

On Nevai's Characterization of Measures with Almost Everywhere Positive Derivative

X. LI AND E. B. SAFF*

*Institute for Constructive Mathematics, Department of Mathematics,
 University of South Florida, Tampa, Florida 33620*

Communicated by Paul Nevai

Received December 1, 1989

It is known that if $d\mu$ is a finite positive Borel measure on the unit circle $\partial\mathcal{A} := \{z \in \mathbb{C} : |z| = 1\}$ with $\mu' > 0$ a.e. in $[0, 2\pi]$, then

$$\lim_{n \rightarrow \infty} \int_0^{2\pi} \left| \frac{|\varphi_n(z)|^2}{|\varphi_{n+l}(z)|^2} - d\theta = 0, \quad z = e^{i\theta} \quad (*)$$

uniformly in $l \geq 0$, where $\{\varphi_n(z)\}_{n=0}^\infty$ is the set of orthonormal polynomials with respect to $d\mu$. Recently, Nevai proved that if (*) holds uniformly in $l \geq 0$, then $\mu' > 0$ a.e. in $[0, 2\pi]$. In this note we establish another characterization of such measures in terms of Turán-type inequalities and we utilize it to give an alternative proof of Nevai's result. © 1990 Academic Press, Inc.

Let $d\mu$ be a finite positive Borel measure on the unit circle $\partial\mathcal{A} := \{z \in \mathbb{C} : |z| = 1\}$ with its support an infinite set. Let $\varphi_n(z) = \varphi_n(d\mu, z) := \kappa_n z^n + \dots \in \mathcal{P}_n$, $\kappa_n > 0$, $n = 0, 1, 2, \dots$, be the n th orthonormal polynomial with respect to $d\mu$; that is,

$$\frac{1}{2\pi} \int_{\partial\mathcal{A}} \varphi_m(z) \overline{\varphi_n(z)} d\mu = \delta_{m,n}, \quad m, n = 0, 1, 2, \dots$$

Rahmanov proved (cf. [9, p. 106])

THEOREM A. *If $\mu' > 0$ a.e. in $[0, 2\pi]$, then*

$$\lim_{n \rightarrow \infty} \frac{z\varphi_n(z)}{\varphi_{n+1}(z)} = 1 \quad (1)$$

uniformly for $|z| \geq 1$

* Research supported, in part, by the National Science Foundation under Grant DMS-881-4026.

In [5] (see also [8]), Máté *et al.* gave a much simpler proof of Theorem A. In doing so, they established (cf. [5, Theorem 3, p. 64])

THEOREM B. *If $\mu' > 0$ a.e. in $[0, 2\pi]$, then*

$$\lim_{l \rightarrow \infty} \int_0^{2\pi} \frac{|\varphi_n(z)|^2}{|\varphi_{n+l}(z)|^2} d\theta = 0, \quad z = e^{i\theta}, \quad (2)$$

uniformly in $l \geq 0$.

On the one hand, Theorem A follows from Theorem B (cf. [5, Theorem 2, p. 64]). On the other hand, as noted in [5, Remark (a), p. 64], Theorem B follows from Theorem A except for the uniformity in l . In fact, recently Nevai (cf. [7, Theorem 1.1, p. 295]) proved that (2) implies $\mu' > 0$ a.e. in $[0, 2\pi]$. In this paper we establish another characterization of measures with positive derivative and we use it to give an alternative proof of the above mentioned result of Nevai.

In [4], the following inequality of Turán played an essential role.

THEOREM C (cf. [6, Corollary 7.5, p. 258]). *If $\mu' > 0$ a.e. in $[0, 2\pi]$, then for every $\delta \in (0, 2\pi]$ there is an $\varepsilon > 0$ such that*

$$\int_E |\varphi_n|^2 d\mu \geq \varepsilon \quad (3)$$

holds for every $n \geq 0$ and every Borel set $E \subset \partial\Delta$ with Lebesgue measure $|E| \geq \delta$.

We are concerned with the following question: Is $\mu' > 0$ a.e. in $[0, 2\pi]$ necessary for (3)? To answer this question, we prove the following refinement of Theorem C which characterizes measures with positive derivative in terms of Turán-type inequalities.

THEOREM 1. *Let $d\mu$ be a finite positive Borel measure on $\partial\Delta$ with infinite support. Then $\mu' > 0$ a.e. in $[0, 2\pi]$ if and only if for every $\delta \in (0, 2\pi]$ and $\rho \in (0, 1)$, there exists an integer $N(\delta, \rho)$ such that*

$$\int_E |\varphi_n|^2 d\mu \geq \rho\delta, \quad n \geq N(\delta, \rho), \quad (4)$$

for all Borel sets $E \subset \partial\Delta$ with $|E| \geq \delta$.

Proof. Let us first assume that $\mu' > 0$ a.e. in $[0, 2\pi]$. Then by [6, Theorem 7.4, p. 257],

$$\lim_{n \rightarrow \infty} \int_E |\varphi_n|^2 d\mu = |E|$$

oof of

uniformly for all Borel sets $E \subset \partial\Delta$. Thus, given $\delta \in (0, 2\pi]$ and $\rho \in (0, 1)$, we can find an integer $N(\delta, \rho)$ such that

$$\left| \int_E |\varphi_n|^2 d\mu - |E| \right| < (1 - \rho)\delta, \quad n \geq N(\delta, \rho),$$

(2)

uniformly for all Borel sets $E \subset \partial\Delta$. Hence, when $n \geq N(\delta, \rho)$,

$$\int_E |\varphi_n|^2 d\mu \geq |E| - (1 - \rho)\delta \geq \rho\delta$$

of [5,
p. 64],
in fact,
 $\mu' > 0$
ion of
proof

for all Borel sets $E \subset \partial\Delta$ with $|E| \geq \delta$, which establishes (4).

Now assume that (4) holds. From the Lebesgue decomposition (cf. [10, p. 21])

$$d\mu = \mu' d\theta + d\mu_s,$$

where $d\mu_s \perp d\theta$. Assume that $d\mu_s$ is concentrated on a Borel set A . Then

$$|A| = 0.$$

$[0, 2\pi]$,

Given $\varepsilon > 0$, take $\delta = 2\pi$ and $\rho \in (0, 1)$ such that $2\pi - \rho\delta = 2\pi(1 - \rho) < \varepsilon$. Then, for $n \geq N(2\pi, \rho)$, we get from (4) with $E = A^c$

measure

$$\int_A |\varphi_n|^2 d\mu = \int_0^{2\pi} |\varphi_n|^2 d\mu - \int_{A^c} |\varphi_n|^2 d\mu \leq 2\pi - \rho\delta < \varepsilon.$$

$[0, 2\pi]$
refine-
ivative

Note that

$$\int_0^{2\pi} |\varphi_n|^2 d\mu_s = \int_A |\varphi_n|^2 d\mu_s = \int_A |\varphi_n|^2 d\mu,$$

infinite
 $[\pi]$ and

and so

$$\int_0^{2\pi} |\varphi_n|^2 d\mu_s < \varepsilon,$$

when $n \geq N(2\pi, \rho)$. Hence,

$$\lim_{n \rightarrow \infty} \int_0^{2\pi} |\varphi_n|^2 d\mu_s = 0. \quad (5)$$

by [6,

Now set

$$B := \{\theta \in [0, 2\pi] : \mu'(\theta) = 0\}$$

If $|B| =: \delta_0 > 0$, then for $n \geq N(\delta_0, \frac{1}{2})$, we have from (4)

$$\frac{1}{2}\delta_0 \leq \int_B |\varphi_n|^2 d\mu = \int_B |\varphi_n|^2 d\mu_s \leq \int_0^{2\pi} |\varphi_n|^2 d\mu_s,$$

which contradicts (5). Therefore $|B| = 0$, i.e., $\mu' > 0$ a.e. in $[0, 2\pi]$. ■

Now we state and prove Nevai's result (cf. [7, Theorem 1.1, p. 295]).

THEOREM 2. *Let $d\mu$ be a finite positive Borel measure on $\partial\Delta$ with infinite support. Then $\mu' > 0$ a.e. in $[0, 2\pi]$ if and only if (2) holds uniformly in $l \geq 0$.*

In the proof of Theorem 2 we make use of the following lemma in [6, Lemma 4.2, p. 248]. It is a consequence of [2, Theorem 5.2.2, p. 198] (also see [1, formula (1.20), p. 7]).

LEMMA 3. *For any $f \in C(\partial\Delta)$,*

$$\lim_{l \rightarrow \infty} \int_0^{2\pi} \frac{f(z)}{|\varphi_n(z)|^2} d\theta = \int_0^{2\pi} f(z) d\mu, \quad z = e^{i\theta}.$$

Proof of Theorem 2. In view of Theorem B, we only need show that (2) implies $\mu' > 0$ a.e. in $[0, 2\pi]$. To do so, it suffices, from Theorem 1 to show that (2) implies (4).

Assume that (2) holds. Let $\delta \in (0, 2\pi]$ and $\rho \in (0, 1)$ be given. We will proceed in three steps.

Step I. We first show that (4) holds uniformly for every set E which is a finite union of open intervals.

Assume that

$$E = \bigcup_{i=1}^m (\alpha_i, \beta_i) \subset \partial\Delta, \quad (\alpha_i, \beta_i) \cap (\alpha_j, \beta_j) = \emptyset, \quad i \neq j, \quad (6)$$

and $|E| \geq \delta$.

It is easy to see that there exists an $f_{E,\rho} \in C(\partial\Delta)$ and $E_1 =: \bigcup_{i=1}^m (\alpha'_i, \beta'_i) \subset E$ such that

$$\chi_{E_1}(z) \leq f_{E,\rho}(z) \leq \chi_E(z), \quad z \in \partial\Delta$$

(where χ_K denotes the characteristic function of the set $K \subset \mathbb{C}$), and

$$|E_1| = \rho^{1/2} |E|$$

By (2), we can find an integer $M(\delta, \rho)$ such that, for $n \geq M(\delta, \rho)$,

$$\int_0^{2\pi} f_{E,\rho}(z) \left| \frac{|\varphi_n(z)|^2}{|\varphi_{n+l}(z)|^2} - 1 \right| d\theta \leq \int_0^{2\pi} \left| \frac{|\varphi_n|^2}{|\varphi_{n+l}|^2} - 1 \right| d\theta < (\rho^{1/2} - \rho)\delta$$

uniformly in $l \geq 0$ and E is of the form given in (6). Thus, for $n \geq M(\delta, \rho)$,

$$\begin{aligned} \int_0^{2\pi} f_{E,\rho} \frac{|\varphi_n|^2}{|\varphi_{n+l}|^2} d\theta &\geq \int_0^{2\pi} f_{E,\rho} d\theta - (\rho^{1/2} - \rho)\delta \\ &\geq \rho^{1/2} \int_0^{2\pi} f_{E,\rho} d\theta \geq \rho^{1/2} \int_E d\theta = \rho |E| \end{aligned}$$

uniformly in $l \geq 0$ and E of the form given in (6) with $|E| \geq \delta$. From Lemma 3, by letting $l \rightarrow \infty$, we get

$$\int_0^{2\pi} f_{E,\rho} |\varphi_n|^2 d\mu \geq \rho |E|, \quad n \geq M(\delta, \rho).$$

Hence,

$$\int_E |\varphi_n|^2 d\mu \geq \int_0^{2\pi} f_{E,\rho} |\varphi_n|^2 d\mu \geq \rho |E|, \quad n \geq M(\delta, \rho),$$

uniformly for E of the form given in (6) with $|E| \geq \delta$.

Step II. Next we show that (4) holds uniformly for every open subset E of $\partial\Delta$ with $|E| \geq \delta$.

We can write such an open set as

$$(6) \quad E = \bigcup_{i=1}^{\infty} (\alpha_i, \beta_i), \quad (\alpha_i, \beta_i) \cap (\alpha_j, \beta_j) = \emptyset, \quad i \neq j,$$

and

$$|E| = \sum_{i=1}^{\infty} (\beta_i - \alpha_i) \geq \delta.$$

Let m be so large that

$$\sum_{i=1}^m (\beta_i - \alpha_i) \geq \rho^{1/2} \delta$$

Then, when $n \geq M(\rho^{1/2}\delta, \rho^{1/2})$, where $M(\cdot, \cdot)$ is given by Step I, we have

$$\int_E |\varphi_n|^2 d\mu \geq \int_{\bigcup_{i=1}^m (\alpha_i, \beta_i)} |\varphi_n|^2 d\mu \geq \rho^{1/2} \cdot \rho^{1/2}\delta = \rho\delta$$

uniformly for any open set $E \subset \partial\Delta$ with $|E| \geq \delta$.

Step III. We complete the proof by showing that E can be any Borel set with $|E| \geq \delta$.

For such a set E we can find, by the outer-regularity of measure $|\varphi_n|^2 d\mu$, an open set $\mathcal{O}_{n,E} \supset E$ such that

$$\rho^{-1/2} \int_E |\varphi_n|^2 d\mu \geq \int_{\mathcal{O}_{n,E}} |\varphi_n|^2 d\mu.$$

Therefore, by Step II and the inequality $|\mathcal{O}_{n,E}| \geq \delta$, we have

$$\int_E |\varphi_n|^2 d\mu \geq \rho^{1/2} \int_{\mathcal{O}_{n,E}} |\varphi_n|^2 d\mu \geq \rho^{1/2} \cdot \rho^{1/2}\delta = \rho\delta,$$

when $n \geq M(\rho^{1/4}\delta, \rho^{1/4})$. ■

Finally, we remark that in [3], Lubinsky gave an explicit expression of a pure jump distribution (i.e., a discrete measure) dv on $\partial\Delta$ such that (cf. [3, formula (2.22), p. 523])

$$\sum_{j=2 \cdot 3^{l-1} + 1}^{2 \cdot 3^l} |\Phi_j(0)|^2 \leq C \left(\frac{\log l}{l} \right),$$

l large enough, where $\Phi_j(z) := (1/\kappa_j)\varphi_j(z)$ and $C > 0$ is a constant independent of l . Consequently,

$$\lim_{n \rightarrow \infty} |\Phi_n(0)| = 0,$$

which is equivalent to (1) (cf., e.g., [5, formula (7), p. 66]). Summarizing these remarks, we can state

PROPOSITION 4. *There exists a finite positive Borel measure that is a pure jump distribution but for which (1) holds.*

REFERENCES

1. G. FREUD, "Orthogonal Polynomials," Pergmon Press, New York, 1971.
2. YA. L. GEROMINUS, "Orthogonal Polynomials," Pergmon, New York, 1960.

- ave
- Borel
- $\int_n |^2 d\mu,$
3. D. S. LUBINSKY, Jump distributions on $[-1, 1]$ whose orthogonal polynomials have leading coefficients with given asymptotic behavior, *Proc. Amer. Math. Soc.* **104** (1988), 516–524.
 4. X. LI, E. B. SAFF, AND Z. SHA, Behavior of best L_p polynomial approximants on the unit interval and on the unit circle, *J. Approx. Theory*, to appear.
 5. A. MÁTÉ, P. NEVAI, AND V. TOTIK, Asymptotics for the ratio of leading coefficients of orthonormal polynomials on the unit circle, *Constr. Approx.* **1** (1985), 63–69.
 6. A. MÁTÉ, P. NEVAI, AND V. TOTIK, Strong and weak convergence of orthogonal polynomials, *Amer. J. Math.* **109** (1987), 239–282.
 7. P. NEVAI, Characterization of measures associated with orthogonal polynomials on the unit circle, *Rocky Mountain J. Math.* **19** (1989), 293–302.
 8. P. NEVAI, Weakly convergent sequences of functions and orthogonal polynomials, *J. Approx. Theory* **65** (1991), in press.
 9. E. A. RAHMANOV, On the asymptotics of the ratio of orthogonal polynomials, II, *Math. USSR-Sb.* **46** (1983), 105–117.
 10. W. RUDIN, “Real and Complex Analysis,” 3rd ed., McGraw–Hill, New York, 1986.
- sion of
at (cf.
- t inde-
- arizing
- a pure