

ABRAHAMSE'S INTERPOLATION THEOREM AND FUCHSIAN GROUPS

MRINAL RAGHUPATHI

ABSTRACT. We generalize Abrahamse's interpolation theorem from the setting of a multiply connected domain to that of a more general Riemann surface. Our main result provides the scalar-valued interpolation theorem for the fixed-point subalgebra of H^∞ associated to the action of a Fuchsian group. We rely on two results from a paper of Forelli. This allows us to prove the interpolation result using duality techniques that parallel Sarason's approach to the interpolation problem for H^∞ . In this process we prove a more general distance formula, very much like Nehari's theorem, and obtain relations between the kernel function for the character automorphic Hardy spaces and the Szegő kernel for the disk. Finally, we examine our interpolation results in the context of the two simplest examples of Fuchsian groups acting on the disk.

1. INTRODUCTION

1.1. **Motivation.** Our objective in this paper is a generalization of Abrahamse's interpolation theorem [1, Theorem 1]. Let $R \subseteq \mathbb{C}$ be a bounded, multiply connected domain of genus g . Let $H^\infty(R)$ be the Banach algebra of bounded holomorphic functions on R . Abrahamse considered a natural class of Hilbert modules $\{H_\lambda^2(R) : \lambda \in \mathbb{T}^g\}$ over the algebra $H^\infty(R)$. Each of the spaces $H_\lambda^2(R)$ is a reproducing kernel Hilbert space with kernel function K^λ . Abrahamse's solution to the Pick interpolation theorem for the region R is the following:

Theorem 1.1 (Abrahamse). *Let $z_1, \dots, z_n \in R$ and $w_1, \dots, w_n \in \mathbb{D}$. Then there exists a holomorphic function $f : R \rightarrow \mathbb{D}$ such that $f(z_j) = w_j$ if and only if the matrices*

$$A_\lambda := [(1 - w_i \overline{w_j}) K^\lambda(z_i, z_j)] \geq 0,$$

for all $\lambda \in \mathbb{T}^g$.

There are several proofs of Abrahamse's theorem in the literature: see Ball [8], for the matrix-valued generalization; Fedorov and Vinnikov [15], for a careful analysis of the scalar-valued problem; and McCullough [22], and McCullough and Paulsen [23], for the operator algebraic viewpoint.

In this paper we take a viewpoint that in some sense predates all the above proofs. Let $p : \mathbb{D} \rightarrow R$ be the universal covering map. Associated to this covering map is

Date: January 19, 2009.

2000 Mathematics Subject Classification. Primary 47A57; Secondary 46E22, 30F35.

Key words and phrases. Hardy spaces, Nevanlinna-Pick interpolation, Distance formulae, Nehari's theorem, Fuchsian groups.

The author wishes to acknowledge the support and guidance of his advisor, Vern I. Paulsen. This work was completed as part of the author's PhD thesis at the University of Houston in August 2008. I would also like to thank the referee for his comments and careful reading of the manuscript.

the *group of deck transformations*, that is, the group Γ of Möbius transformations such that $p \circ \gamma = p$. Note that the map $p^* : H^\infty(R) \rightarrow H^\infty(\mathbb{D})$ given by $p^*(f) = f \circ p$ is an isometric embedding of $H^\infty(R)$, and the range of p^* is precisely the fixed-point subalgebra for the action of Γ , that is, the set of functions $f \in H^\infty(\mathbb{D})$ such that $f \circ \gamma = f$ for all $\gamma \in \Gamma$. Therefore, function theoretic problems on the domain R can be lifted to the disk. This idea, known as *uniformization*, is a fundamental tool in the theory of Riemann surfaces [14]. If Γ is the group of deck transformations associated to p , then Γ is a discrete subgroup of the group of all Möbius transformations on the disk. In addition, Γ is torsion-free, Γ acts without fixed points and the action is properly discontinuous. Further, the quotient space $\mathbb{D} \backslash \Gamma$ is a Riemann surface that can be identified with R , and $\Gamma \cong \pi_1(R)$, the fundamental group of R .

A discrete group of Möbius transformations is called a *Fuchsian group*. Let Γ be a Fuchsian group. We will consider the analogue of Abrahamse's theorem for H_Γ^∞ , the associated fixed-point subalgebra of H^∞ . From comments in the previous paragraph we see that this includes the study of the interpolation problem on a Riemann surface.

Our approach in this paper owes a great deal to the elegant results obtained by Forelli [17]. Forelli initiated the study of function theory in these fixed-point algebras, and in the special case where Γ is a group of deck transformations associated to a covering of a finite bordered Riemann surface, he constructs a bounded projection from $P : H^\infty \rightarrow H_\Gamma^\infty$ that is also a bimodule map over H_Γ^∞ . The projection is obtained by modifying the natural conditional expectation from H^∞ onto H_Γ^∞ . An explicit formula for the conditional expectation was obtained by Earle and Marden [12] and this was generalized by Ball [8].

Another approach to function theory on multiply connected domains through the use of holomorphic vector bundles was initiated by Abrahamse and Douglas [4] using the results in [17]. More recently Ball and Vinnikov [9] have obtained a great deal of information about Hardy spaces on finite bordered Riemann surfaces by further developing the theory of holomorphic vector bundles and through the viewpoint of complex algebraic curves. Fedorov and Vinnikov [15] carried out a deep analysis of the scalar-valued interpolation problem for finite open Riemann surfaces using complex analytic methods.

We study the interpolation problem from the perspective of classical Hardy space theory with an additional element: the action of a Fuchsian group. Although we do not provide a complete geometric or topological description of which Riemann surfaces are included in our result, our results will apply to any Riemann surface that has a finitely generated fundamental group. This includes the class of finite bordered Riemann surfaces and finite open Riemann surfaces. In addition, we allow the possibility that the group Γ may act with fixed points or possess torsion. We will need neither the full strength of Forelli's results nor the extensions obtained in [8] and [12]. We also do not require a knowledge of function theory on multiply connected domains or the theory of holomorphic vector bundles. However, our proof does rest on two lemmas in [17].

1.2. Notation. We will denote the open unit disk in the complex plane by \mathbb{D} . The circle will be denoted \mathbb{T} and the m -torus will be denoted \mathbb{T}^m . By an *automorphism* we mean a Möbius transformation, that is, a holomorphic map of \mathbb{D} onto \mathbb{D} with a holomorphic inverse. An automorphism is the composition of a simple Blaschke

factor and a rotation, that is, an automorphism ϕ is of the form

$$\phi(z) = \lambda \frac{a - z}{1 - \bar{a}z},$$

where $a \in \mathbb{D}$ and $\lambda \in \mathbb{T}$. The derivative ϕ is easily computed to be

$$\phi'(z) = \lambda \frac{|a|^2 - 1}{(1 - \bar{a}z)^2}.$$

Hence, given an automorphism ϕ , the constants a and λ are determined by the equations

$$a = \phi^{-1}(0), \lambda = (|a|^2 - 1)^{-1} \phi'(0).$$

The set of all automorphisms of the disk is a group under the composition of maps. This group is naturally identified with $PSL(2, \mathbb{R})$.

We will use the term *automorphism group* to mean a subgroup of the group of all automorphisms of the disk and we will usually denote an automorphism group by Γ . An automorphism of the disk is a well-defined holomorphic map on the open set $\{z : |z| < |a|^{-1}\} \supseteq \mathbb{D} \cup \mathbb{T}$. Therefore, the action of Γ on the disk extends to the boundary circle and defines an action of Γ on the circle \mathbb{T} .

Given a group Γ of automorphisms and a point $w \in \mathbb{D}$ we denote the stabilizer of the point w by $\text{Fix}(\Gamma, w)$, that is,

$$\text{Fix}(\Gamma, w) := \{\gamma \in \Gamma : \gamma(w) = w\}.$$

We denote the orbit of the point w under the action of Γ by $\Gamma(w)$, that is,

$$\Gamma(w) := \{\gamma(w) : \gamma \in \Gamma\}.$$

We will call two groups Γ_1 and Γ_2 *conjugate* if and only if there exists an automorphism ϕ such that $\phi\Gamma_1 = \Gamma_2\phi$. We denote by $[\Gamma, \Gamma]$ the commutator subgroup of Γ .

The group Γ acts, by composition, on the Lebesgue spaces $L^p := L^p(\mathbb{T})$, for $1 \leq p \leq \infty$, with respect to normalized Lebesgue measure μ . Since a Möbius transformation is analytic this action restricts to the Hardy spaces H^p .

If $X \subseteq L^p$ we denote by $[X]_p$ the smallest closed subspace of L^p that contains X . If $p = 2$, then we write $[X]_2 = [X]$. When $p = \infty$ we take closures with respect to the weak* topology.

The character group, or dual group, of Γ is the set of continuous homomorphisms from the group Γ into the circle \mathbb{T} . Let $X \subseteq L^p$ be a set that is closed under composition by elements of Γ . If $\sigma \in \hat{\Gamma}$, then we define

$$X_{\sigma} := \{f \in X : f \circ \gamma = \sigma(\gamma)f\}.$$

We call this the *character automorphic space associated to σ* . We call elements of X_{σ} *character automorphic*.

1.3. Outline of our work. Since our focus is on function theoretic results, we are interested in the cases where the group Γ fixes some nonconstant function in $H_{\mathbb{F}}^{\infty}$, that is, when $H_{\mathbb{F}}^{\infty}$ is nontrivial. If $f \in H_{\mathbb{F}}^{\infty}$ is nontrivial, then we can, after subtracting a constant, assume that $f(0) = 0$. Therefore, $f(\gamma(0)) = 0$ for all $\gamma \in \Gamma$ and so the points in the orbit of the origin $\Gamma(0)$ must satisfy the Blaschke condition

$$(1) \quad \sum_{\zeta \in \Gamma(0)} (1 - |\zeta|) < \infty.$$

In this case $\sum_{\gamma \in \Gamma} (1 - |\gamma(w)|)$ converges for any point $w \in \mathbb{D}$. An elementary calculation shows that the sum in (1) converges if and only if the Poincaré series $\sum_{\gamma \in \Gamma} |\gamma'(w)|$ converges for any point $w \in \mathbb{D}$. Automorphism groups for which either, and hence both, of these series converges are said to be of *convergence type*; see Tsuji [30, Theorem XI.3]. It is straightforward to see that if Γ is a group of convergence type, then Γ is Fuchsian.

If the series in (1) converges, then we can form the Blaschke product with zero set $\Gamma(0)$ and this Blaschke product is character automorphic. In Section 2, we will use these Blaschke products to construct orthonormal bases for the character spaces H_σ^2 . In Section 3 we review some of the results from Forelli [17]. In particular, we revisit the defect space N . We assume, as did Forelli, that the space N is finite-dimensional. We recall some duality results for the Hardy spaces H_Γ^p . These duality results are a generalization of Forelli's original result [17, Lemma 3], and the result of Earle and Marden [12, Proposition 5]. The proof we provide is purely functional analytic.

We call a group Γ *admissible* if and only if Γ is a Fuchsian group of convergence type for which the defect space is finite dimensional. In Section 4, we look more closely at the reproducing kernel function for the space H_σ^2 and show how it is related to the Szegő kernel. These results imply that H_σ^∞ is dense in the space H_σ^p . In section 5, we examine more closely the structure of the operator algebra $H_\Gamma^\infty \subset B(H_\Gamma^2)$. The main results of our paper are proved in Section 6. In particular, we prove a distance formula, Theorem 6.1, which implies the following analogue of Nehari's theorem:

Theorem 1.2. *Let Γ be Fuchsian group of convergence type with finite-dimensional defect space N , that is, let Γ be admissible. Let $f \in L_\Gamma^\infty$. The distance of f from H_Γ^∞ is given by*

$$\|f + H_\Gamma^\infty\| = \sup_{\sigma \in \hat{\Gamma}} \|(I - P_{H_\sigma^2})M_f P_{H_\sigma^2}\|.$$

By specializing to the case where $f \in H_\Gamma^\infty$, this distance formula leads to a generalization of Abrahamse's theorem.

Theorem 1.3. *Let Γ be admissible. Let $z_1, \dots, z_n \in \mathbb{D}$ and $w_1, \dots, w_n \in \mathbb{C}$. There exists a function $f \in H_\Gamma^\infty$ with $\|f\|_\infty \leq 1$ such that $f(z_j) = w_j$ if and only if*

$$A_\sigma := [(1 - w_i \bar{w}_j)K^\sigma(z_i, z_j)]_{i,j=1}^n \geq 0$$

for all $\sigma \in \hat{\Gamma}$.

Finally, in Section 7 we illustrate our ideas in the context of the two simplest examples of Fuchsian groups.

2. FACTORIZATION AND BLASCHKE PRODUCTS

2.1. Factorization results. The first step in our work is to establish the analogues of the two main factorization theorems for Hardy spaces: the inner-outer factorization and the Riesz factorization. The inner-outer factorization appears in one form or another in Abrahamse [2, Theorem 1.12], Hasumi [18, Lemma 1], and Voichick and Zalcman [32]. A version of the Riesz factorization is in Abrahamse [1, Lemma 5]. As we will make frequent use of these results we state and give a short proof of them.

Let Γ be an automorphism group. Recall that an element $f \in L^p$ is called *character automorphic* if and only if there exists a character $\sigma \in \hat{\Gamma}$ such that $f \circ \gamma = \sigma(\gamma)f$ for all $\gamma \in \Gamma$. A function $f \in L^p$ is said to be *modulus automorphic* if and only if $|f| \in L^p_\Gamma$. The absolute value of a character automorphic element of L^p is modulus automorphic. A function $u \in H^p$ is called *outer* if and only if the closed linear span in H^p of $\{z^n u : n \geq 0\}$ is all of H^p . The following result shows that a modulus automorphic outer function is character automorphic.

Proposition 2.1. *If $u \in H^p$ is an outer function and $|u| \in L^p_\Gamma$, then there exists $\sigma \in \hat{\Gamma}$ such that $u \in H^p_\sigma$.*

Proof. If u is outer, then $u \circ \gamma$ is outer for all $\gamma \in \Gamma$, since composition by γ is continuous and invertible on H^p . Two outer function u and v have equal modulus if and only if there exists a scalar $\lambda \in \mathbb{T}$ such that $u = \lambda v$, see [13, Corollary 6.23]. We have $|u \circ \gamma| = |u| \circ \gamma = |u|$ and so there exists $\sigma(\gamma) \in \mathbb{T}$ such that $u \circ \gamma = \sigma(\gamma)u$. We need to show that $\sigma \in \hat{\Gamma}$. If $\gamma_1, \gamma_2 \in \Gamma$, then

$$\sigma(\gamma_1 \gamma_2)u = u \circ (\gamma_1 \gamma_2) = (u \circ \gamma_1) \circ \gamma_2 = \sigma(\gamma_1)(u \circ \gamma_2) = \sigma(\gamma_1)\sigma(\gamma_2)u.$$

Since u is nonzero, σ is a character. \square

We can now state the two factorization theorems that we need. We refer the reader to [19, Chapter 4.4] for a statement and proof of these factorization results in the classical situation.

Proposition 2.2. *Let $\sigma \in \hat{\Gamma}$, let $f \in H^p_\sigma$, and let $f = Bsu$ be the canonical factorization of f into a Blaschke product B , a singular inner function s , and an outer function u . Then there exist characters $\sigma_1, \sigma_2, \sigma_3 \in \hat{\Gamma}$ such that $\sigma = \sigma_1 \sigma_2 \sigma_3$, $B \in H^\infty_{\sigma_1}$, $u \in H^1_{\sigma_2}$ and $s \in H^\infty_{\sigma_3}$.*

Proof. Note that if $f(z) = 0$, then $f(\gamma(z)) = 0$ and so the zeros of f are made up of the union of countably many disjoint orbits. The Blaschke product B vanishes precisely on the zero set of f . Since γ permutes the orbit of a point, we see that $B \circ \gamma$ also vanishes on the zero set of f and so $B \circ \gamma = BC$ where C is inner. A similar argument shows that $B \circ \gamma^{-1} = BD$ with D an inner function. We have,

$$B = B \circ \gamma \circ \gamma^{-1} = (BC) \circ \gamma^{-1} = (B \circ \gamma^{-1})(C \circ \gamma^{-1}) = BD(C \circ \gamma^{-1}).$$

Since H^∞ has no zero divisors we get that $D(C \circ \gamma^{-1}) = 1$ and so $\bar{D} = C \circ \gamma^{-1} \in H^\infty$. This shows that D and C are constant and so $B \circ \gamma = \sigma_1(\gamma)B$ for some scalar $\sigma_1(\gamma) \in \mathbb{T}$. We now proceed as in the proof of Proposition 2.1 to check that $\sigma_1 \in \hat{\Gamma}$.

Since $|f| = |u| \in L^1_\Gamma$, we see by Proposition 2.1 that $u \in H^1_{\sigma_3}$ for some character σ_3 . It now follows from the uniqueness of the factorization that $s \in H^\infty_{\sigma_2}$, where $\sigma_2 = \sigma(\sigma_1 \sigma_3)^{-1}$. \square

One consequence of the proof of Proposition 2.2 is the following: if the zero set of the Blaschke product B is the orbit $\Gamma(z)$, then $B \in H^\infty_\sigma$ for some $\sigma \in \hat{\Gamma}$. This fact has been observed many times in the literature [1] and we will make this formal later on in this section.

Proposition 2.3 (Riesz factorization). *Let Γ be an automorphism group and let $\sigma \in \hat{\Gamma}$. If $f \in H^1_\sigma$, then there exists characters $\sigma_1, \sigma_2 \in \hat{\Gamma}$ with $\sigma = \sigma_1 \sigma_2^2$, an inner function $\phi \in H^\infty_{\sigma_1}$, and an outer function $u \in H^2_{\sigma_2}$ such that $f = \phi u^2$.*

Proof. It is well known [19, Theorem 19] that f has a factorization of the form ϕu^2 , where ϕ is inner and $u \in H^2$ is outer. Note that $|f|^{1/2} = |u| \in L^2_\Gamma$ and so by Proposition 2.1, $u \in H^2_{\sigma_2}$ for some $\sigma_2 \in \hat{\Gamma}$. It follows easily that $\phi \in H^\infty_{\sigma_1}$ with $\sigma_1 = \sigma\sigma_2^{-2}$. \square

2.2. Character automorphic Blaschke products. If Γ is of convergence type, then there is an associated class of character-automorphic Blaschke products. These Blaschke products occur naturally when we carry out the inner-outer factorization for elements of H^1_Γ . For us, the primary use of these Blaschke products is in constructing an orthonormal basis for H^2_σ . It is easy to check that a Blaschke product B is character automorphic if and only if its zero set is an at most countable union of orbits. We begin by analyzing the convergence of the Blaschke sum (1).

Proposition 2.4 ([21, Theorem 2.3.5]). *Let Γ be a group of automorphisms. If $\text{Fix}(\Gamma, 0)$ is finite, then $\text{Fix}(\Gamma, 0)$ is cyclic.*

Proposition 2.5. *Let Γ be a group of automorphisms of the disk. Assume that the algebra H^∞_Γ is nontrivial, that is, contains a nonconstant function. The following are true:*

- (1) *For every $w \in \mathbb{D}$ the stabilizer $\text{Fix}(\Gamma, w)$ is a finite, cyclic group.*
- (2) *For every $w \in \mathbb{D}$, the Blaschke sum $\sum_{\gamma \in \Gamma} (1 - |\gamma(w)|)$ converges.*

Proof. Let $w \in \mathbb{D}$. Choose an automorphism ϕ that maps w to 0. Let $\Gamma' = \phi\Gamma\phi^{-1}$ and note that $\text{Fix}(\Gamma', 0) = \text{Fix}(\Gamma, w)$ and $\Gamma'(0) = \phi(\Gamma(w))$. Also note that the spaces $H^\infty_{\Gamma'}$ and H^∞_Γ are isometrically isomorphic. Therefore, it is enough to consider the case $w = 0$.

- (1) Every element $\gamma \in \text{Fix}(\Gamma, 0)$ fixes the origin. Hence, γ is a rotation of the disk and there exists a constant $\lambda \in \mathbb{T}$ such that $\gamma(z) = \lambda z$. Let f be a nonconstant function in H^∞_Γ and let a_k , $k \neq 0$, be a nonzero Fourier coefficient of f . For every $\gamma \in \text{Fix}(\Gamma, 0)$ we have $f(\gamma(z)) = f(z)$ and so $\lambda^k a_k = a_k$. This yields, $\lambda^k = 1$ and so $\text{Fix}(\Gamma, 0)$ is finite. The claim about $\text{Fix}(\Gamma, 0)$ being cyclic follows from Proposition 2.4.
- (2) If $f \in H^\infty_\Gamma$ is nonconstant, then by subtracting a constant we may assume that f is nonzero and vanishes at 0. Since $f \circ \gamma = f$ we see that $f(\gamma(0)) = f(0) = 0$ for all $\gamma \in \Gamma$ and so $\sum_{\zeta \in \Gamma(0)} (1 - |\zeta|) < \infty$. If $\alpha \in \Gamma$, then the cardinality of $\text{Fix}(\Gamma, 0)$ is equal to the cardinality of $\text{Fix}(\Gamma, \alpha(0))$. In fact, the two stabilizers are isomorphic via the map $\gamma \mapsto \alpha\gamma\alpha^{-1}$. Therefore,

$$\sum_{\gamma \in \Gamma} (1 - |\gamma(0)|) = |\text{Fix}(\Gamma, 0)| \sum_{\zeta \in \Gamma(0)} (1 - |\zeta|) < \infty.$$

\square

Note that if the series $\sum_{\gamma \in \Gamma} (1 - |\gamma(0)|)$ does converge, then the argument above shows that $\text{Fix}(\Gamma, 0)$ is finite and Γ is discrete.

If the stabilizer $\text{Fix}(\Gamma, w)$ is finite, and the series $\sum_{\gamma \in \Gamma} (1 - |\gamma(w)|)$ converges, then we define two Blaschke products B_w and $B_{\Gamma, w}$ that arise naturally. First consider the Blaschke product B_w whose zero set is $\Gamma(w)$. We call B_w the *Blaschke product for the orbit* $\Gamma(w)$. Let $m = |\text{Fix}(\Gamma, w)|$ and define $B_{\Gamma, w} := B_w^m$. We call B_Γ the *Blaschke product associated to the pair* $(\Gamma, 0)$. If it is the case that only the identity map fixes w , then $m = 1$ and $B_{\Gamma, w} = B_w$. If $\gamma \in \Gamma$, then $B_w \circ \gamma$ is a

Blaschke product whose zero set is the same as the zero set of B_w . It follows, just as in Proposition 2.2, that $B_w \circ \gamma = \sigma_w(\gamma)B_w$ for some character σ_w . We call σ_w the *character associated to the pair* (Γ, w) . We will most often be interested in the case where $w = 0$, in which case we denote $B_{\Gamma,0}$ by B_Γ .

We now construct an orthonormal basis for H_σ^2 from the Blaschke product B_0 and the kernel functions for the spaces H_σ^2 . Our purpose in the next few lemmas is to show that it is enough to construct an orthonormal basis in the case where H_σ^2 contains at least one function that does not vanish at the origin.

Lemma 2.6. *Let $\gamma \in \Gamma$ be a generator of the cyclic group $\text{Fix}(\Gamma, 0)$ and let $\gamma(z) = \lambda z$, where $\lambda \in \mathbb{T}$. Let $\gamma_n(0)$ be an enumeration of the distinct nonzero elements of $\Gamma(0)$. Then $B_0 \circ \gamma = \lambda B_0$ and $B_\Gamma \circ \gamma = B_\Gamma$.*

Proof. The Blaschke product B_0 can be written

$$B_0(z) = z \prod_{n=1}^N \frac{|\gamma_n(0)|}{\gamma_n(0)} \frac{\gamma_n(0) - z}{1 - \overline{\gamma_n(0)}z}.$$

Hence,

$$B_0(\lambda z) = (\lambda z) \prod_{n=1}^N \frac{|\gamma_n(0)|}{\gamma_n(0)} \frac{\gamma_n(0) - \lambda z}{1 - \overline{\gamma_n(0)}\lambda z} = \lambda z \prod_{n=1}^N \frac{|\bar{\lambda}\gamma_n(0)|}{\bar{\lambda}\gamma_n(0)} \frac{\bar{\lambda}\gamma_n(0) - z}{1 - \overline{\gamma_n(0)}\lambda z}.$$

We have, $\bar{\lambda}\gamma_n(0) = \gamma^{-1}(\gamma_n(0))$ and so $\bar{\lambda}\gamma_n(0)$ is another enumeration of the nonzero points in the orbit $\Gamma(0)$. Hence, $B_0(\lambda z) = \lambda B_0(z)$. If $|\text{Fix}(\Gamma, 0)| = m$, then $\gamma(z) = e^{\frac{2\pi i}{m}}(z)$ is a generator for $\text{Fix}(\Gamma, 0)$ and so $B_\Gamma(e^{\frac{2\pi i}{m}}z) = (e^{\frac{2\pi i}{m}}B_0(z))^m = B_\Gamma$ for all $z \in \mathbb{D}$. \square

Lemma 2.7. *Let $f \in H_\sigma^1$ be a character automorphic function. Let $m = |\text{Fix}(\Gamma, 0)|$ and let $\rho(z) = e^{2\pi i/m}z$ be the generator of $\text{Fix}(\Gamma, 0)$. Then $\sigma(\rho) = e^{2\pi ir/m}$, $0 \leq r \leq m-1$. If $r \neq 0$, then $f(0) = 0$. If $r = 0$ and $f(0) = 0$, then $B_\Gamma|f$.*

Proof. Let $\Gamma_1 = \langle \rho \rangle$ and let $\omega = e^{2\pi i/m}$. Since ρ has order m , we see that there exists an integer r , $0 \leq r \leq m-1$, such that $\sigma(\rho) = \omega^r$. Since $f \circ \rho = \sigma(\rho)f = \omega^r f$, we see that the Fourier expansion of f has the form $f = \sum_{j=0}^{\infty} a_{mj+r} z^{mj+r}$. If $r \neq 0$, then $f(0) = 0$. On the other hand if $r = 0$, then $z^m|(f - f(0))$. Hence, $B_\Gamma|(f - f(0))$. \square

The Hardy space H^2 is a reproducing kernel Hilbert space, and the kernel function for H^2 is the Szegő kernel

$$K^S(z, w) = \frac{1}{1 - \bar{w}z}.$$

If \mathcal{M} is a subspace of H^2 , then \mathcal{M} is also a reproducing kernel Hilbert space and we denote its kernel function $K^\mathcal{M}$. We denote the kernel function for H_σ^2 by K^σ and the kernel function for H_Γ^2 by K^Γ . Given a point $w \in \mathbb{D}$ and a kernel function K we denote by k_w , the kernel function at the point w , that is, k_w is the unique element of \mathcal{M} such that $\langle f, k_w \rangle = f(w)$ for all $f \in \mathcal{M}$. It follows that $\|k_w\|^2 = K(w, w)$. If $k_w \neq 0$, then we define the normalized kernel function by $\kappa_w = K(w, w)^{-1/2}k_w$. Otherwise, we set $\kappa_w = 0$. Recall that Γ is of convergence type if the sum in (1) converges.

Proposition 2.8. *Let Γ be a Fuchsian group of convergence type. Let $m = |\text{Fix}(\Gamma, 0)|$. Assume that $\chi \in \hat{\Gamma}$ and that there exists a function $f \in H_\sigma^2$ with $f(0) \neq 0$. For $n \geq 0$, let $\chi_n = \chi\sigma_0^{-mn}$, let $H_n^2 = H_{\chi_n}^2$ and let $\kappa_0^{(n)}$ denote the normalized kernel function for H_n^2 . Then the nonzero elements of the set*

$$\mathcal{E}_\chi := \{B_\Gamma^n \kappa_0^{(n)} : n \geq 0\},$$

is an orthonormal basis for H_χ^2 .

Proof. Since B_0 is an inner function and $\kappa_0^{(n)}$ is the normalized kernel function we see that the nonzero elements of \mathcal{E}_χ have norm 1. Since $B_0(0) = 0$, we see that the nonzero elements of \mathcal{E}_χ are orthogonal.

If ρ is the generator of $\text{Fix}(\Gamma, 0)$, then $\chi(\rho) = 1$, by Lemma 2.7. Combining this with Lemma 2.6 we see that $\chi_n(\rho) = 1$ for $n \geq 1$.

Now let $f \in H_\chi^2$ with $f \perp \mathcal{E}_\chi$. Note that $\kappa_0^{(n)} = 0$ implies that every element of H_n^2 vanishes at the origin. Since $f \perp \kappa_0^{(0)}$, we see that $f(0) = 0$ and so $B_\Gamma|f$, by Lemma 2.7. Therefore, $f = B_\Gamma f_1$ and by composing with elements of Γ we see that $f_1 \in H_{\chi\sigma_0^{-m}}^2 = H_{\chi_1}^2$. If $k_0^{(1)} = 0$, then every element of H_1^2 vanishes at the origin and so $B_\Gamma|f_1$. If $k_0^{(1)} \neq 0$, then the fact that $f \perp B_\Gamma \kappa_0^{(1)}$ gives $f_1(0) = 0$ and $B_\Gamma|f_1$. Repeating this process we see that B_Γ^n divides f for all $n \geq 0$. Hence, $f = 0$. \square

3. THE FORELLI PROJECTION

3.1. The conditional expectation. To begin this section we review some ideas from the work of Forelli [17]. Let Γ be an automorphism group. Let \mathfrak{M} denote the collection of Lebesgue measurable subsets of \mathbb{T} . For $E \in \mathfrak{M}$, define E to be Γ -invariant if and only if $\mu(E \Delta \gamma^{-1}(E)) = 0$ for every $\gamma \in \Gamma$. Forelli proved the following fact: the collection of Γ -invariant sets \mathfrak{M}_Γ is a sub-sigma-algebra of \mathfrak{M} , and the subspace $L^p(\mathbb{T}, \mathfrak{M}_\Gamma, \mu)$ is equal to L_Γ^p .

It follows, from standard arguments in probability theory, that there is the unique conditional expectation $\Psi_p : L^p(\mathbb{T}, \mathfrak{M}, \mu) \rightarrow L^p(\mathbb{T}, \mathfrak{M}_\Gamma, \mu)$. The conditional expectation Ψ_p is weak*-continuous for $1 < p \leq \infty$, and $\Psi_p|L^q = \Psi_q$ for $p \leq q$. Therefore, we omit the subscript p and denote Ψ_p by Ψ . Note that the map Ψ is a projection ($\Psi^2 = \Psi$), and is selfadjoint ($\Psi^* = \Psi$) when $p = 2$. The map Ψ has the property that

$$\int_E \Psi(f) d\mu = \int_E f d\mu,$$

for $E \in \mathfrak{M}_\Gamma$. Given $f \in L^p(\mathbb{T}, \mathfrak{M}, d\mu)$ and $g \in L^q(\mathbb{T}, \mathfrak{M}_\Gamma, d\mu)$ we have

$$\Psi(fg) = \Psi(f)g.$$

In the classical setting, the space L^2 decomposes as a direct sum of the Hardy space H^2 and the space $\overline{H_0^2}$, that is, the set of complex conjugates of functions in H^2 that have mean value 0. In our setting we must contend with the fact that the direct sum $H_\Gamma^2 \oplus \overline{H_{0,\Gamma}^2}$ may not be all of L_Γ^2 . The orthogonal complement of $H_\Gamma^2 \oplus \overline{H_{0,\Gamma}^2}$ in L_Γ^2 is called the *defect space*. If Γ is the group of deck transformations that arise from a covering map of a g -holed region, then the defect space is a g -dimensional subspace. Forelli captures the defect space without reference to multiply connected domains. We now outline these ideas, as they are central to our work. We also feel that our presentation is sufficiently different from [17] to merit inclusion.

Let $k_z \in H^2$ denote the Szegő kernel for the Hardy space. To begin with let $f \in L^2_\Gamma$ and let $f = g + \bar{h} + c$, where $g, h \in H^2_0$ and $c = \int f dm$. While the functions g, h are not Γ -invariant, they are “nearly” so. We have

$$g + \bar{h} + c = f = f \circ \gamma = g \circ \gamma + \overline{h \circ \gamma} + c.$$

Rearranging this last equation gives

$$g \circ \gamma - g = \overline{h - h \circ \gamma}.$$

Since $\gamma \in \Gamma$ is analytic we see that $g \circ \gamma$ and $h \circ \gamma$ are both in H^2 and so $g \circ \gamma - g = \overline{h - h \circ \gamma} = c(f, \gamma)$, where $c(f, \gamma)$ is a constant that depends on f and γ . The constant is easily computed by integrating

$$c(f, \gamma) = \int g \circ \gamma - g = g(\gamma(0)) - g(0) = \langle g, k_{\gamma(0)} \rangle.$$

Similarly, $c(f, \gamma) = \langle \bar{h}, \overline{k_{\gamma(0)}} \rangle$. Combining these two equations, we get

$$\begin{aligned} 2c(f, \gamma) &= \langle g + \bar{h}, k_{\gamma(0)} - \overline{k_{\gamma(0)}} \rangle \\ &= \langle \Psi(f), k_{\gamma(0)} - \overline{k_{\gamma(0)}} \rangle = \langle f, \Psi(k_{\gamma(0)} - \overline{k_{\gamma(0)}}) \rangle. \end{aligned}$$

If we set $v_\gamma = -i\Psi(k_{\gamma(0)} - \overline{k_{\gamma(0)}})$, then v_γ is a positive element of L^∞_Γ with the property that

$$\int f v_\gamma = 2ic(f, \gamma)$$

for all $\gamma \in \Gamma$. Note that $-i(k_{\gamma(0)} - \overline{k_{\gamma(0)}}) = P^*_{\gamma(0)}$, the conjugate of the Poisson kernel at the point $\gamma(0)$. Following Forelli we define the *defect space* N by

$$N := \text{span}\{v_\gamma : \gamma \in \Gamma\}.$$

If $f \in L^2_\Gamma$ is orthogonal to v_γ , then $g \circ \gamma - g = 0 = h - h \circ \gamma$. Hence, g and $h \in H^2_{0,\Gamma}$ and we get that $L^2_\Gamma = H^2_\Gamma \oplus \overline{H^2_{0,\Gamma}} \oplus [N]$.

If $\gamma_1, \gamma_2 \in \Gamma$, then

$$\begin{aligned} c(f, \gamma_1 \circ \gamma_2) &= g \circ (\gamma_1 \circ \gamma_2) - g = g \circ (\gamma_1 \circ \gamma_2) - g \circ \gamma_2 + g \circ \gamma_2 - g \\ &= (g \circ \gamma_1 - g) \circ \gamma_2 + (g \circ \gamma_2 - g) = c(f, \gamma_1) + c(f, \gamma_2) \end{aligned}$$

It follows that

$$\int f v_{\gamma_1 \circ \gamma_2} = 2ic(f, \gamma_1 \circ \gamma_2) = 2i(c(f, \gamma_1) + c(f, \gamma_2)) = \int f(v_{\gamma_1} + v_{\gamma_2}).$$

Since this is true for all $f \in L^2_\Gamma$ we get $v_{\gamma_1 \circ \gamma_2} = v_{\gamma_1} + v_{\gamma_2}$.

What all of this shows is that the map $\gamma \mapsto v_\gamma$ is a homomorphism from Γ into the additive group N . This homomorphism must factor through the commutator subgroup $[\Gamma, \Gamma]$ to give a homomorphism from $\Gamma/[\Gamma, \Gamma]$ into N .

Given a set of generators $\{\gamma_s : s \in S\} \subseteq \Gamma/[\Gamma, \Gamma]$ the vectors $v_s := v_{\gamma_s}$, for $s \in S$, span the space N . If the group Γ is finitely generated, then the space N is finite dimensional and the dimension of N is smaller than the minimal number of generators of $\Gamma/[\Gamma, \Gamma]$. Note that if an element $\gamma \in \Gamma$ has finite order, say m , then $mv_\gamma = v_{\gamma^m} = 0$ and so $v_\gamma = 0$. If $\Gamma/[\Gamma, \Gamma]$ is generated by elements of finite order, then N is trivial.

In the case where Γ is the group of deck transformations associated with a universal covering map of a multiply connected domain, Forelli showed that the dimension of N is equal to the minimal number of generators of $\Gamma/[\Gamma, \Gamma]$. Many results follow

from this equality, including a corona theorem for H_Γ^∞ , the key stepping stone being the construction of a bounded projection $P : H^\infty \rightarrow H_\Gamma^\infty$. We do not assume, and do not need, this stronger condition.

We are now in a position to state what kind of groups we will deal with in this paper. We will call a group Γ *admissible* if and only if Γ is of convergence type and the defect space N is finite dimensional.

These are natural, nontrivial conditions from a function theory point of view. The assumption about N means that $H_\Gamma^p + N$ is a closed subspace of L^p for all $1 \leq p \leq \infty$. It is useful to keep in mind that $N \subset L_\Gamma^\infty \subseteq L_\Gamma^p$ for all $1 \leq p \leq \infty$. Note that when $q > 1$, the spaces H_Γ^q and $H_\Gamma^q + N$ are also weak* closed. This is a simple consequence of the fact that weak* limits preserve point values and the fact that N is assumed finite dimensional.

3.2. Duality and density results. Since duality arguments will play a central role in our interpolation results we would like to gather some results on the duality between the different H_Γ^p spaces. The results contained in the next proposition have appeared in various forms in the literature [6, Theorem 11.1], [12, Proposition 5], and [17, Lemma 3]. We provide, mostly for the sake of completeness, an elementary proof that depends only on standard duality arguments and the expectation Ψ .

Proposition 3.1. *Let Γ be an admissible group. The image of H^p under the conditional expectation Ψ is $H_\Gamma^p + N$. For $1 \leq p < \infty$, the dual of L_Γ^p can be identified with L_Γ^q , where $q = \frac{p}{p-1}$. In this identification the following are true:*

- (1) $(H_\Gamma^p)^\perp = H_{0,\Gamma}^q + N$
- (2) $(H_\Gamma^q)^\perp = H_{0,\Gamma}^p + N$
- (3) $(H_\Gamma^q + N)^\perp = H_{0,\Gamma}^p$
- (4) $(H_\Gamma^p + N)^\perp = H_{0,\Gamma}^q$

Proof. The statement about L_Γ^p spaces follows from standard facts about the L^p spaces of a probability measure.

For $p = 2$, we have seen already that $L_\Gamma^2 = H_\Gamma^2 \oplus \overline{H_{0,\Gamma}^2} \oplus N$ and the above results are valid. We have

$$\int \Psi(h)g = \int hg = 0,$$

for all $h \in H^2$, $g \in H_{0,\Gamma}^2$, and so $\Psi(H^2) = L_\Gamma^2 \ominus \overline{H_{0,\Gamma}^2} = H_\Gamma^2 \oplus N$.

Next consider the case $1 \leq p < \infty$. We will show that $H_\Gamma^p = (H_{0,\Gamma}^q + N)^\perp$ and the remaining results will follow either by duality or by a similar argument. Note that the spaces H_Γ^q and $H^q + N$ are weak* closed in L^q .

We first show that $\Psi(H^p) = H_\Gamma^p + N$. Let $f \in L_\Gamma^q$. Note that $f \in \Psi(H^p)^\perp$ if and only if $\int f\Psi(g) = \int fg = 0$ for all $g \in H^p$ if and only if $f \in H_0^q \cap L_\Gamma^q = H_{0,\Gamma}^q$. Hence, $\Psi(H^p)^\perp = H_{0,\Gamma}^q$.

Let $h \in H_\Gamma^p$, $v \in N$ and $g \in H_{0,\Gamma}^q$. Since $v_\gamma \in L_\Gamma^\infty$ and g is analytic we get

$$\int gv_\gamma = \int g\Psi(k_{\gamma(0)} - \overline{k_{\gamma(0)}}) = \int g(k_{\gamma(0)} - \overline{k_{\gamma(0)}}) = g(\gamma(0)) - g(0) = 0.$$

It follows that

$$\int g(h + v) = \int gh + \int gv = 0,$$

which gives $H_\Gamma^p + N \subseteq (H_{0,\Gamma}^q)^\perp$. This also yields

$$\Psi(H^p) = (\Psi(H^p)^\perp)^\perp = (H_{0,\Gamma}^q)^\perp \supseteq H_\Gamma^p + N.$$

Let $g \in (H_\Gamma^p + N)^\perp$ and let $f \in H^p$. We will show that

$$\int gf = \int g\Psi(f) = 0,$$

which will establish the fact that $(H_\Gamma^p + N)^\perp \subseteq \Psi(H^p)^\perp$. We know that $\int g(h+v) = 0$ for all $h \in H_\Gamma^p$ and $v \in N$. Let $f_n \in H^\infty$ and suppose that $f_n \rightarrow f$ in the L^p norm. Since $f_n \in H^2$, we can write $\Psi(f_n) = h_n + v_n$, where $h_n \in H_\Gamma^2$ and $v_n \in N$. However, $v_n \in L_\Gamma^\infty$ and so $h_n \in H_\Gamma^\infty \subseteq H_\Gamma^p$. Therefore,

$$\int g(h_n + v_n) = \int g\Psi(f_n) = 0$$

for all n , since $g \in \Psi(H^p)^\perp$. Hence,

$$\int g\Psi(f) = \lim_{n \rightarrow \infty} \int g\Psi(f_n) = 0.$$

We have established that $\Psi(H^p)^\perp = H_{0,\Gamma}^q$ and that $\Psi(H^p)^\perp \supseteq (H_\Gamma^p + N)^\perp$. This combined with the fact that $\Psi(H^p) \supseteq H_\Gamma^p + N$ shows us that $\Psi(H^p) = H_\Gamma^p + N$ and $(H_\Gamma^p + N)^\perp = H_{0,\Gamma}^q$. \square

Proposition 3.2. For $1 \leq p < \infty$, $[H_\Gamma^\infty]_p = H_\Gamma^p$.

Proof. Consider the case $p \geq 2$. Let $f \in H_\Gamma^p$ and let $f_n \in H^\infty$ converge to f in L^p . Since the L^p norm dominates the L^2 norm we see that $f_n \rightarrow f$ in L^2 . If we project f_n onto $\Psi(f_n)$, then there exists $g_n \in H_\Gamma^\infty$ and $v_n \in N$ such that $\Psi(f_n) = g_n + v_n$ and $g_n + v_n \rightarrow f$ in the L^p norm and also in the L^2 norm. In particular, $\|v_n\|_2 \rightarrow 0$. However, on the finite dimensional space N , the L^2 norm and the L^p norm are equivalent and so $\|v_n\|_p \rightarrow 0$. It follows that $g_n \rightarrow f$ in L^p .

Let $1 \leq p < 2$, view H_Γ^∞ as a subspace of H_Γ^2 , and let $(H_\Gamma^\infty)^\perp$ denote the annihilator of H_Γ^∞ in L_Γ^q , where $2 \leq q$ and $q^{-1} + p^{-1} = 1$. We have

$$\begin{aligned} (H_\Gamma^\infty)^\perp &= \left\{ f \in L_\Gamma^q : \int fg = 0 \text{ for all } g \in H_\Gamma^\infty \right\} \\ &\subseteq \left\{ f \in L_\Gamma^2 : \int fg = 0 \text{ for all } g \in H_\Gamma^\infty \right\} \\ &= L_\Gamma^2 \ominus [\overline{H_\Gamma^\infty}] = L_\Gamma^2 \ominus \overline{H_\Gamma^2} = H_{0,\Gamma}^2 \oplus N. \end{aligned}$$

Hence, $(H_\Gamma^\infty)^\perp \subseteq L_\Gamma^q \cap (H_{0,\Gamma}^2 + N) = H_{0,\Gamma}^q + N$, since $N \subseteq L_\Gamma^\infty \subseteq L_\Gamma^q$. The reverse inclusion follows from the inclusions $H_{0,\Gamma}^q + N \subseteq H_{0,\Gamma}^2 \oplus N$ and $H_\Gamma^\infty \subseteq H_\Gamma^2$, and the result for $p = 2$. We have shown that $(H_\Gamma^\infty)^\perp = H_{0,\Gamma}^q + N$. It follows from Proposition 3.1 that $[H_\Gamma^\infty]_p = ((H_\Gamma^\infty)^\perp)^\perp = (H_{0,\Gamma}^q + N)^\perp = H_\Gamma^p$. \square

We now prove a density result that is central to the proof of our distance formula and interpolation theorem. Recall that a function $u \in H^2$ is called *outer* if and only if $[H^\infty u] = H^2$. If $\sigma \in \hat{\Gamma}$ is a character and $u \in H_\sigma^2$ is an outer function, then $[H_\Gamma^\infty u] \subseteq H_\sigma^2$. The purpose of the next two results is to prove the reverse containment. Let us temporarily adopt the following notation: if $\|u\|_2 = 1$, then

we call u normalized; and let the L^2 norm induced by the measure $|u|^2 dm$ be denoted $\|\cdot\|_{2,u}$.

Lemma 3.3. *Let Γ be an admissible group, let $u \in H^2$ be an outer function and let N be finite dimensional. Then $[H_\Gamma^\infty u] \cap [Nu] = \{0\}$.*

Proof. We can assume that u is normalized. Since N is finite dimensional $[Nu] = Nu$. Let $f \in [H_\Gamma^\infty u] \cap Nu$ and note that $f \in H^2$. Since u is an outer function, there exists a sequence $f_n \in H_\Gamma^\infty$ such that $\lim_{n \rightarrow \infty} \|f - f_n u\|_2 = 0$. Also, there exists $v \in N$ such that $f = vu$. There exists a sequence $h_n \in H^\infty$ such that $\lim_{n \rightarrow \infty} \|h_n u - 1\|_2 = 0$. We have, $v = \lim_{n \rightarrow \infty} v u h_n = \lim_{n \rightarrow \infty} f h_n \in H_\Gamma^2$. This implies $v \in N \cap H_\Gamma^2 = \{0\}$ and so $f = 0$. \square

We are now in a position to prove our main lemma about cyclic subspaces.

Proposition 3.4. *Let Γ be an admissible group, let $\sigma \in \hat{\Gamma}$ and let $u \in H_\sigma^2$ be an outer function. The cyclic subspace generated by H_Γ^∞ and u is equal to H_σ^2 .*

Proof. One inclusion is straightforward, namely $[H_\Gamma^\infty u] \subseteq H_\sigma^2$.

For the converse, let $f \in H_\sigma^2 \ominus [H_\Gamma^\infty u]$. We will show that $f = 0$. Since u is an outer function, there exists a sequence of functions $f_n \in H^\infty$ such that $\lim_{n \rightarrow \infty} \|f_n u - f\|_2 = 0$. Let Ψ be the conditional expectation from L^1 onto L_Γ^1 . Note that $|u|^2, f\bar{u}$ and $|f|^2 \in L_\Gamma^1$. Using the fact that $\int \Psi(f)g = \int fg$ for $f \in L^\infty$ and $g \in L_\Gamma^1$ we get

$$\begin{aligned} \|\Psi(f_n)u - f\|_2^2 &= \int |\Psi(f_n)|^2 |u|^2 - \Psi(f_n)u\bar{f} - \overline{\Psi(f_n)u}f + |f|^2 dm \\ &= \int \overline{f_n}\Psi(f_n) |u|^2 - \Psi(f_n)u\bar{f} - \overline{f_n}u\bar{f} + |f|^2 dm \\ &= \int (\Psi(f_n)u - f)\overline{(f_n u - f)} dm = \langle \Psi(f_n)u - f, f_n u - f \rangle \\ &\leq \|\Psi(f_n)u - f\|_2 \|f_n u - f\|_2. \end{aligned}$$

Hence, $\lim_{n \rightarrow \infty} \|\Psi(f_n)u - f\|_2 \leq \lim_{n \rightarrow \infty} \|f_n u - f\|_2 = 0$.

By Proposition 3.1 we can write $\Psi(f_n) = g_n + v_n$, where $g_n \in H_\Gamma^\infty$ and $v_n \in N$. Hence, $f \in [H_\Gamma^\infty u] + [Nu]$. Since N is finite dimensional, by Lemma 3.3, we can define an equivalent norm on $[H_\Gamma^\infty u] + [Nu]$ by $\|h + w\|' = \|h\|_2 + \|w\|_2$. Since $f \perp g_n u$, it follows that $\lim_{n \rightarrow \infty} \|v_n u - f\|' = 0$, and $f \in [Nu] \cap [H_\Gamma^\infty u] = \{0\}$. \square

4. REPRODUCING KERNEL FUNCTIONS FOR CHARACTER SPACES

The purpose of the results in this section is to establish connections between the reproducing kernel K^σ and the Szegö kernel K^S . As a corollary to these results we will see that H_σ^∞ is dense in H_σ^1 .

Lemma 4.1. *If Γ is an admissible group, then $(H_\Gamma^1 + N) \cap (\overline{H_\Gamma^1} + N) = N + \mathbb{C}$.*

Proof. Let $f \in (H_\Gamma^1 + N) \cap (\overline{H_\Gamma^1} + N)$. We can write $f = g + v = \bar{h} + w$, where $g, h \in H_{0,\Gamma}^1$ and $v, w \in N$. We have $g - \bar{h} = w - v \in N$. By Proposition 3.2 we can choose sequences $g_n, h_n \in H_\Gamma^\infty$ such that $g_n \rightarrow g$, $h_n \rightarrow h$ in L_Γ^1 . Note that $g_n - \bar{h}_n \in H_\Gamma^2 \oplus \overline{H_\Gamma^2}$ and so $g_n - \bar{h}_n \perp N$. Therefore, $\|g - \bar{h}\|_1 = 0$ which yields $g \in H_\Gamma^1 \cap \overline{H_\Gamma^1}$. Hence, g is constant and $f \in N + \mathbb{C}$. \square

Proposition 4.2. *Let Γ be an admissible group, let $\sigma \in \hat{\Gamma}$, let $\mathcal{N} := H_\sigma^2 \ominus [H_{0,\Gamma}^\infty H_\sigma^2]$. If $f \in \mathcal{N}$, then $|f|^2 \in N + \mathbb{C}$. In particular, the kernel function at the origin k_0^σ is bounded.*

Proof. Let $f \in \mathcal{N}$ and let $h \in H_{0,\Gamma}^\infty$. We have $f \perp hf$, that is,

$$\int |f|^2 h = 0,$$

for all $h \in H_{0,\Gamma}^\infty$. By taking the complex conjugate we also get

$$\int |f|^2 \bar{h} = 0,$$

and so $|f|^2 \in N + \mathbb{C}$, by Lemma 4.1. Since $N \subseteq L_\Gamma^\infty$, this implies that $|f|$ is bounded.

To prove the claim about the kernel function, note that $k_0^\sigma \perp [H_{0,\Gamma}^\infty H_\sigma^2]$. \square

Corollary 4.3. *Let Γ be an admissible group. If $\sigma \in \hat{\Gamma}$, then H_σ^∞ is dense in H_σ^p .*

Proof. It is enough to prove the result in the case $p = 1$. Recall that, in Proposition 2.8, we had constructed an orthonormal basis for H_σ^2 . Each of the elements of this basis was the product of a Blaschke product and a normalized kernel function at the origin for some character space. These are bounded by Proposition 4.2 and we see that H_σ^∞ is dense in H_σ^2 . If $f \in H_\sigma^1$, then by Riesz factorization (see Proposition 2.3) there exists $g \in H_{\sigma_1}^2$ and $h \in H_{\sigma\sigma_1^{-1}}^2$ such that $f = gh$. Choose a sequence $g_n \in H_{\sigma_1}^\infty$ and $h_n \in H_{\sigma\sigma_1^{-1}}^\infty$ such that $\|g_n - g\|_2 \rightarrow 0$ and $\|h_n - h\|_2 \rightarrow 0$ in H^2 . It follows that f is the limit of $g_n h_n \in H_\sigma^\infty$ and so H_σ^∞ is dense in H_σ^1 . \square

We now show that the kernel function for H_χ^2 is bounded.

Corollary 4.4. *Let Γ be an admissible group, let $w \in \mathbb{D}$ and let B_Γ be the Blaschke product associated to the pair $(\Gamma, 0)$. Then there exists a constant C , which depends only on the group Γ , such that*

$$(2) \quad \|k_w^\chi\|_\infty \leq C^2(1 - |B_\Gamma(w)|)^{-1}$$

for every $\chi \in \hat{\Gamma}$.

Proof. Since $N + \mathbb{C}$ is finite dimensional we know that there is a constant C such that $\|f\|_\infty \leq C\|f\|_2$ for all $f \in N + \mathbb{C}$. Hence, $\|k_0^\chi\|_\infty \leq C\|k_0^\chi\|_2$ and $\|\kappa_0^\chi\|_\infty \leq C$. Note that this bound is independent of the character χ .

From Proposition 2.8 we have an orthonormal basis for the space H_χ^2 consisting of the nonzero elements of the set $\mathcal{E}_\chi = \{B_\Gamma^n \kappa_0^{(n)} : n \geq 0\}$. For ease of notation let $B = B_\Gamma$ and $f_n = \kappa_0^{(n)}$. We have

$$K^\chi(z, w) = \sum_{n=0}^{\infty} B^n(z) f_n(z) \overline{B^n(w) f_n(w)}.$$

Taking the absolute value and using the fact that $\|\kappa_0^\chi\|_\infty \leq C$ we get

$$|K^\chi(z, w)| \leq C^2 \sum_{n=0}^{\infty} |B(z)|^n |B(w)|^n \leq \frac{C^2}{1 - |B(z)B(w)|}.$$

Therefore, $|k_w^\chi(z)| \leq C^2(1 - |B(z)B(w)|)^{-1} \leq C^2(1 - |B(w)|)^{-1}$. Taking the supremum over $z \in \mathbb{D}$ yields the inequality in (2). \square

5. H_{Γ}^{∞} AS AN OPERATOR ALGEBRA

5.1. Commutants and multiplier algebras. In this section we provide some basic information about the operator algebra structure of H_{Γ}^{∞} when viewed as the multiplier algebra of H_{σ}^2 .

Before we proceed to our results we recall the definition of a multiplier algebra. Given a reproducing kernel Hilbert space (RKHS) \mathcal{H} on the set X with kernel function K , the multiplier algebra $\text{mult}(\mathcal{H})$ is the set of functions $f : X \rightarrow \mathbb{C}$ such that $fh \in \mathcal{H}$ for all $h \in \mathcal{H}$. The product here is taken pointwise. If $f \in \text{mult}(\mathcal{H})$, then the operator $M_f : \mathcal{H} \rightarrow \mathcal{H}$ defined by $M_f(h) = fh$ is a bounded linear transformation. The multiplier norm of f is defined by $\|f\|_{\text{mult}} = \|M_f\|$. It is not difficult to show that $M_f^*k_x = \overline{f(x)}k_x$. From this we see that f is a bounded function and $\|f\|_{\text{mult}} \geq \|f\|_{\infty}$. We begin by identifying the multiplier algebra of H_{σ}^2 , where $\sigma \in \hat{\Gamma}$.

Proposition 5.1. *If $\sigma \in \hat{\Gamma}$, then the multiplier algebra of H_{σ}^2 is H_{Γ}^{∞} .*

Proof. If $f \in H_{\Gamma}^{\infty}$ and $g \in H_{\sigma}^2$, then it is easily checked that $fg \in H_{\sigma}^2$.

For the converse we can assume that no inner function divides H_{σ}^2 . Hence, $k_w^{\sigma} \neq 0$ for all $w \in \mathbb{D}$ and we see that a multiplier f of H_{σ}^2 is a bounded function. Now let $h \in H_{\sigma}^2$, with $h \neq 0$, and let $\gamma \in \Gamma$. We have $(fh) \circ \gamma = \sigma(\gamma)fh$. On the other hand, $(fh) \circ \gamma = (f \circ \gamma)(h \circ \gamma) = \sigma(\gamma)(f \circ \gamma)h$. Comparing these two equations we see that $f \circ \gamma = f$, that is, $f \in H_{\Gamma}^{\infty}$.

It is a well-known fact [24, Theorem 6.3] that $\|f\|_{\text{mult}} \geq \|f\|_{\infty}$. On the other hand, given $f \in H_{\Gamma}^{\infty}$ and $h \in H_{\sigma}^2$, we have

$$\|M_f h\|_2^2 = \|fh\|_2^2 = \int |fh|^2 dm \leq \|f\|_{\infty}^2 \|h\|_2^2.$$

Hence, $\|f\|_{\text{mult}} = \|M_f\| \leq \|f\|_{\infty}$. Combining these inequalities we get $\|f\|_{\text{mult}} = \|f\|_{\infty}$. \square

We see from Proposition 5.1 that the multiplier algebra of H_{σ}^2 is H_{Γ}^{∞} and that the multiplier norm is the usual supremum norm. My considering the map $f \mapsto M_f$ we have identified H_{Γ}^{∞} as an algebra of operators on the space H_{σ}^2 , that is, we have identified H_{Γ}^{∞} with a unital, weak* closed subalgebra of $B(H_{\sigma}^2)$.

Invariant subspaces will now play a central role in our results. We recall the notion of *reflexivity* for an operator algebra $\mathcal{A} \subseteq B(\mathcal{H})$. We denote by $\text{Lat}(\mathcal{A})$ the lattice of invariant subspaces of \mathcal{A} , that is, the set of projections P such that $PAP = AP$ for all $A \in \mathcal{A}$. An algebra $\mathcal{A} \subseteq B(\mathcal{H})$ is called *reflexive* if and only if $\text{Alg}(\text{Lat}(\mathcal{A})) = \mathcal{A}$.

If \mathcal{L} is the lattice in $B(H^2)$ of subspaces of the form ϕH^2 , where ϕ ranges over the set of inner functions, then $\text{Alg}(\text{Lat}(\mathcal{L})) = H^{\infty}$ and so H^{∞} is reflexive. We now prove a more general fact that implies that H_{Γ}^{∞} is reflexive as a subalgebra of H_{σ}^2 .

Proposition 5.2. *Let \mathcal{H} be a reproducing kernel Hilbert space. If \mathcal{A} is the multiplier algebra of \mathcal{H} , then \mathcal{A} is a reflexive operator algebra in $B(\mathcal{H})$.*

Proof. Suppose that $T \in B(\mathcal{H})$ and $T(\mathcal{M}) \subseteq \mathcal{M}$ for all $\mathcal{M} \in \text{Lat}(\mathcal{A})$. The subspace spanned by the kernel function k_x is invariant for \mathcal{A}^* and so $T^*k_x \in \text{span}\{k_x\}$. Hence, there exists a constant $\phi(x) \in \mathbb{C}$ such that $T^*k_x = \phi(x)k_x$ and consequently $T = M_{\bar{\phi}}$. \square

Our next proposition implies that H^∞_Γ is equal to its own commutant when viewed as a subalgebra of $B(H^2_\Gamma)$.

Proposition 5.3. *Let \mathcal{M} be a subspace of H^2 such that $1 \in \mathcal{M}$ and let \mathcal{A} be the multiplier algebra of this subspace. If $[\mathcal{A}] = \mathcal{M}$, then $\mathcal{A}' = \mathcal{A}$, when \mathcal{A} is represented as multiplication operators on \mathcal{M} .*

Proof. Let $T \in \mathcal{A}'$ and set $T(1) = h$. Consider the action of T^* on the kernel function $k_x \in \mathcal{M}$. Let $g \in \mathcal{A}$ and compute

$$\begin{aligned} \langle T^*k_x, g \rangle &= \langle k_x, TM_g1 \rangle = \langle k_x, M_gT(1) \rangle = \langle k_x, gh \rangle \\ &= \overline{g(x)h(x)} = \left\langle \overline{h(x)}k_x, g \right\rangle \end{aligned}$$

and so $T^*k_x = \overline{h(x)}k_x$. This forces $h \in \mathcal{A}$ and $T = M_h$. \square

If $\gamma : \mathbb{D} \rightarrow \mathbb{D}$ is an automorphism, then $C_\gamma : H^2 \rightarrow H^2$ denotes the composition operator defined by $C_\gamma(f) = f \circ \gamma$. An easy calculation,

$$\langle f, C_\gamma^*k_z \rangle = \langle C_\gamma f, k_z \rangle = f(\gamma(z)) = \langle f, k_{\gamma(z)} \rangle,$$

shows that $C_\gamma^*k_z = k_{\gamma(z)}$. Given a Fuchsian group Γ , let us denote by $\text{Alg}(S, \Gamma)$ the smallest subalgebra of $B(H^2)$ that contains the unilateral shift S and the group of composition operators $\{C_\gamma : \gamma \in \Gamma\}$.

If $h \in H^2$, $f \in H^\infty$ and $\gamma \in \Gamma$, then

$$M_f C_\gamma(h) = f(h \circ \gamma) = (f \circ \gamma^{-1})(h \circ \gamma) = ((f \circ \gamma^{-1})h) \circ \gamma = C_\gamma M_{f \circ \gamma^{-1}}(h).$$

Hence, $M_{f \circ \gamma^{-1}} = C_\gamma^{-1} M_f C_\gamma$. This shows that the action of Γ on H^∞ is implemented by a similarity. The fixed point space for this action is the algebra H^∞_Γ which is the set of $f \in H^\infty$ such that $M_f C_\gamma = C_\gamma M_f$ for all $\gamma \in \Gamma$. We now give the most basic connection between $\text{Alg}(S, \Gamma)$ and H^∞_Γ .

Proposition 5.4. *The commutant of $\text{Alg}(S, \Gamma)$ in $B(H^2)$ is H^∞_Γ .*

Proof. If $T \in \text{Alg}(S, \Gamma)'$, then $TS = ST$ which forces $T = M_f \in H^\infty$. Since $M_f C_\gamma = C_\gamma M_f$, for all $\gamma \in \Gamma$, we get $f \in H^\infty_\Gamma$. \square

5.2. Inner multipliers of the defect space. While the space N does not contain analytic functions, it was shown by Forelli [17, Lemma 7] that it was possible to multiply N into H^2 by an inner function. The following observation about the invariant subspaces of $\text{Alg}(S, \Gamma)$ will prove useful.

Proposition 5.5. *The invariant subspaces of $\text{Alg}(S, \Gamma)$ are of the form ϕH^2 , where ϕ is a character-automorphic inner function.*

Proof. If \mathcal{M} is invariant for $\text{Alg}(S, \Gamma)$, then \mathcal{M} is shift invariant. It follows that $\mathcal{M} = \phi H^2$ for some inner function ϕ . Since \mathcal{M} is invariant under C_γ we get that $(\phi \circ \gamma)H^2 = \phi H^2$. It follows, from the uniqueness statement in Beurling's theorem [19, Chapter 4.4], that $\phi \circ \gamma = \sigma(\gamma)\phi$ where $\sigma(\gamma) \in \mathbb{T}$. If $\gamma_1, \gamma_2 \in \Gamma$, then

$$\sigma(\gamma_1 \gamma_2)\phi = \phi \circ (\gamma_1 \gamma_2) = (\phi \circ \gamma_1) \circ \gamma_2 = (\sigma(\gamma_1)\phi) \circ \gamma_2 = \sigma(\gamma_1)\sigma(\gamma_2)\phi.$$

Hence, $\sigma \in \hat{\Gamma}$. \square

Our next proposition shows that there are no finite-dimensional subalgebras of H^∞ , aside from the constants. We will need this fact for the special case where the subalgebra under consideration is H^∞_Γ .

Proposition 5.6. *If \mathcal{A} is a unital subalgebra of H^∞ that contains nonconstant functions, then \mathcal{A} is infinite dimensional.*

Proof. Assume to the contrary that \mathcal{A} is n -dimensional. Let $f \in \mathcal{A}$ be nonconstant. By subtracting $f(0)$, we may assume that $f(0) = 0$ and $f \neq 0$. The elements $1, f, f^2, \dots, f^n$ must be linearly dependent and so there exists a_0, \dots, a_n such that $\sum_{j=0}^n a_j f^j = 0$. Evaluating at $z = 0$, we get $a_0 = 0$ and so $f(a_1 + a_2 f + \dots + a_n f^{n-1}) = 0$. Since $f \neq 0$, $a_1 + a_2 f + \dots + a_n f^{n-1} = 0$. Repeating the above argument yields $a_1 = \dots = a_n = 0$, a contradiction. \square

As noted by Forelli, the subspace $H_\Gamma^2 + N$ is invariant for H_Γ^∞ . To see this pick $h \in H_\Gamma^2 + N$ and choose $k \in H^2$ such that $\Psi(k) = h$. If $f \in \overline{H_{0,\Gamma}^2}$ and $g \in H_\Gamma^\infty$, then

$$\int f \overline{gh} = \int f \overline{g \Psi(k)} = \int f \overline{gk} = 0.$$

Hence, $H_\Gamma^\infty(H_\Gamma^2 + N) \subseteq H_\Gamma^2 + N$. This shows that N is semi-invariant for H_Γ^∞ and so the compression of H_Γ^∞ to N is a homomorphism of H_Γ^∞ . Since H_Γ^∞ is infinite dimensional and N is finite dimensional we see that this homomorphism has a nontrivial kernel. Hence, there exists a function $f \in H_\Gamma^\infty$ such that $M_f(N) \perp N$.

While the functions in N are not analytic, they can be multiplied into H^2 by a character-automorphic inner function. Let \mathcal{N} be the set of functions $f \in H^2$ such that $fN \subset H^2$. The subspace \mathcal{N} is closed and invariant for $\text{Alg}(S, \Gamma)$. Hence, the subspace $\mathcal{N} = \phi H^2$ for some character-automorphic inner function ϕ , by Proposition 5.5. This means that $\phi N \subseteq H^\infty$ with ϕ character automorphic. This is the proof of [17, Lemma 7]

We point out that there is a related way to obtain this last fact. Let \mathcal{M} denote the smallest shift invariant subspace of L^2 that contains $H_\Gamma^2 + N$. Clearly $\mathcal{M} = [H^\infty(H_\Gamma^2 + N)]$. By the Helson-Lowdenslager theorem [19, Theorem 18], the subspace is either of the form $\chi_E L^2$ or of the form ϕH^2 where ϕ is unimodular. Suppose that $\mathcal{M} = \chi_E L^2$ and note that $\chi_E = 1$, since $1 \in \mathcal{M}$. If $\mathcal{M} = L^2$, then $\Psi(\mathcal{M}) = L_\Gamma^2$. A typical element of \mathcal{M} can be approximated by sums of elements of the form fg where $f \in H^\infty$ and $g \in H_\Gamma^2 + N$. Note that $\Psi(fg) = \Psi(f)g$, which shows

$$\begin{aligned} L_\Gamma^2 &= \Psi(\mathcal{M}) \subseteq [\{\Psi(fg) : f \in H^\infty, g \in H_\Gamma^2 + N\}] \\ &= [\{\Psi(f)g : f \in H^\infty, g \in H_\Gamma^2 + N\}] \\ &= [\{fg : f \in H_\Gamma^\infty + N, g \in H_\Gamma^2 + N\}] \\ &\subseteq H_\Gamma^2 + N + N.N, \end{aligned}$$

which is impossible since N is finite dimensional and $\overline{H_{0,\Gamma}^\infty}$ is not. Hence, $\mathcal{M} = \psi H^2$ for a unimodular character-automorphic function ψ . Since $1 \in H_\Gamma^2 + N \subseteq \psi H^2$, we get $1 = \psi \phi$ for an inner function ϕ and so $\overline{\psi} = \phi$ is inner. Hence, $\phi(H_\Gamma^2 + N) \subseteq H^2$. This ability to multiply N into H^∞ with a character-automorphic inner function is central to our proof of Theorem 6.1 and so we record this fact.

Lemma 5.7 (Forelli). *There exists a character-automorphic inner function ϕ such that $\phi N \subseteq H^\infty$.*

Using Lemma 5.7 we get the following lemma:

Lemma 5.8. *If $w \in \mathbb{D}$ and Γ is an admissible Fuchsian group, then there exists a character-automorphic inner function ϕ_w such that $\phi_w \overline{k_w^\Gamma} \in H^\infty$.*

Proof. We can assume that H_Γ^∞ is nontrivial, otherwise $k_w^\Gamma = 1$. Let $\mathcal{N} = \{f \in H^2 : f \overline{k_w^\Gamma} \in H^2\}$. The subspace \mathcal{N} is closed subspace and invariant for $\text{Alg}(S, \Gamma)$. Since H_Γ^∞ is nontrivial, we can find $f \in H_\Gamma^\infty$ such that $f \neq 0$ and $f(w) = 0$. Let $g \in H_\Gamma^2$ and note that $\langle f \overline{k_w^\Gamma}, \overline{g} \rangle = \langle fg, k_w^\Gamma \rangle = f(w)g(w) = 0$. Hence, $f \overline{k_w^\Gamma} \in H_{0, \Gamma}^2 + N$. By Lemma 5.7, there exists an inner function ϕ such that $\phi f \overline{k_w^\Gamma} \in H^2$. It follows that the space \mathcal{N} is nontrivial and, by Beurling's theorem, there exists an inner function ϕ_w such that $\mathcal{N} = \phi_w H^2$. It follows from Proposition 5.5 that ϕ_w is character automorphic and $\phi_w \overline{k_w^\Gamma} \in H^2$. Since $k_w^\Gamma \in H^\infty$, it follows that $\phi_w \overline{k_w^\Gamma} \in H^\infty$. \square

6. A GENERALIZATION OF ABRAHAMSE'S THEOREM

We now come to a formula for the distance of an element in L_Γ^∞ from the algebra H_Γ^∞ . Actually we will prove a slightly stronger statement. Let $z_1, \dots, z_n \in \mathbb{D}$ and let \mathcal{I} denote the ideal of functions in H_Γ^∞ that vanish at the n points $z_1, \dots, z_n \in \mathbb{D}$. Let \mathcal{K}_σ denote the span of the kernel functions $k_{z_1}^\sigma, \dots, k_{z_n}^\sigma$ and let

$$\mathcal{N}_\sigma = H_\sigma^2 \ominus \mathcal{K}_\sigma = \{f \in H_\sigma^2 : f(z_1) = \dots = f(z_n) = 0\}.$$

By combining Proposition 3.4 with [26, Lemma 4.6] we get that $\mathcal{N}_\sigma = [\mathcal{I}u]$ for any outer function $u \in H_\sigma^2$.

Our first result in this section is a distance formula. The formula is an analogue of Nehari's theorem, which relates the distance of $f \in L^\infty$ from H^∞ to the norm of the Hankel operator with symbol f .

The formula is also related to the Arveson distance formula. To understand this connection we need the notion of hyper-reflexivity [7]. If $A \in \mathcal{A}$, $P \in \text{Lat}(\mathcal{A})$ and $T \in B(\mathcal{H})$, then

$$\|T + A\| \geq \|(I - P)(T + A)P\| = \|(I - P)TP + (I - P)AP\| = \|(I - P)TP\|.$$

From this we see that $\|T + \mathcal{A}\| \geq \sup_{P \in \text{Lat}(\mathcal{A})} \|(I - P)TP\|$. The algebra \mathcal{A} is *hyper-reflexive* if and only if the reverse inequality holds up to a constant, that is, there exists a constant C such that $\|T + \mathcal{A}\| \leq C \sup_{P \in \text{Lat}(\mathcal{A})} \|(I - P)TP\|$ for all $T \in B(\mathcal{H})$.

With these comments in mind we now come to central result.

Theorem 6.1. *Let Γ be an admissible group and let $f \in L_\Gamma^\infty$. The distance of f from \mathcal{I} is given by*

$$(3) \quad \|f + \mathcal{I}\| = \sup_{\sigma \in \Gamma} \|(I - P_{\mathcal{N}_\sigma})M_f P_{H_\sigma^2}\|.$$

Here I denotes the identity in $B(L^2)$ and the orthogonal projections are in $B(L^2)$.

Proof. Let $k_{z_j}^\Gamma$ be the kernel function at the point z_j for the space H_Γ^2 . By duality

$$(4) \quad \|f + \mathcal{I}\| = \sup \left| \int fg \right|$$

where $g \in \mathcal{I}_\perp$ and $\|g\|_1 \leq 1$. Since $\mathcal{I} \subseteq H_\Gamma^\infty$, we see that $\mathcal{I}_\perp \supseteq H_\Gamma^1 + N$. Since the functions in \mathcal{I} vanish at the points $z_1, \dots, z_n \in \mathbb{D}$, we also have $\overline{\mathcal{K}_\Gamma} \subseteq \mathcal{I}_\perp$. An application of Proposition 3.1 shows that $\mathcal{I}_\perp = H_{\Gamma, 0}^1 + N + \overline{\mathcal{K}_\Gamma}$. By Lemma 5.8

there exist character-automorphic inner functions ϕ_1, \dots, ϕ_n such that $\phi_j \overline{k_{z_j}^\Gamma} \in H^2$. Let ϕ_0 be the inner function, as in Lemma 5.7, such that $\phi_0 N \subseteq H^\infty$. Let $\phi = \phi_0 \dots \phi_n$ and note that ϕ is character automorphic. We have $\phi g \in H^1$ and, by Proposition 2.3, there exists a character $\sigma \in \hat{\Gamma}$ and an outer function $u \in H_\sigma^2$ such that $|g| = |u|^2$. Rewriting the expression in (4) we get

$$\|f + \mathcal{I}\| = \sup \left| \int f u \bar{v} \right|$$

where u is an outer function in H_σ^2 and $\bar{v}u = g$. Since $g \in \mathcal{I}_\perp$ we see that $\int gh = 0$ for all $h \in \mathcal{I}$. Therefore, $\langle hu, v \rangle = 0$ and so $v \in L^2 \ominus [\mathcal{I}u] = L^2 \ominus \mathcal{N}_\sigma$. Hence,

$$\left| \int f u \bar{v} \right| = |\langle fu, v \rangle| \leq \|(I - P_{\mathcal{N}_\sigma})M_f P_{H_\sigma^2}\|.$$

We now prove the reverse inequality. Let $g \in \mathcal{I}$ and $f \in L_\Gamma^\infty$. If $h \in H_\sigma^2$, then $gh \in \mathcal{N}_\sigma$ and so $(I - P_{\mathcal{N}_\sigma})M_g P_{H_\sigma^2} = 0$. We have,

$$\begin{aligned} \|f + g\| &= \|M_{f+g}\| \geq \|(I - P_{\mathcal{N}_\sigma})M_{f+g}P_{H_\sigma^2}\| \\ &= \|(I - P_{\mathcal{N}_\sigma})M_f P_{H_\sigma^2} + (I - P_{\mathcal{N}_\sigma})M_g P_{H_\sigma^2}\| \\ &= \|(I - P_{\mathcal{N}_\sigma})M_f P_{H_\sigma^2}\| \end{aligned}$$

□

Our generalization of Abrahamse's theorem now follows from this distance formula. The theorem applies to a fairly broad class of Riemann surfaces. In particular, the theorem applies to quotients of the disk \mathbb{D} by an action of a finitely generated Fuchsian group. In addition, if the action of the group has no fixed points, then the quotient is a Riemann surface with finitely generated fundamental group. Therefore, our results apply to the case of any finite bordered Riemann surface [15, 9]. We also point out that the proof, and the lemmas on which it depends, does not require knowledge of function theory on multiply connected domains.

Theorem 6.2. *Let Γ be an admissible Fuchsian group. Let $z_1, \dots, z_n \in \mathbb{D}$ and $w_1, \dots, w_n \in \mathbb{C}$. There exists a function $f \in H_\Gamma^\infty$ with $\|f\|_\infty \leq 1$ such that $f(z_j) = w_j$ if and only if the matrices*

$$(5) \quad A_\sigma := [(1 - w_i \bar{w}_j)K^\sigma(z_i, z_j)]_{i,j=1}^n \geq 0$$

for all $\sigma \in \hat{\Gamma}$.

Proof. If $f \in H_\Gamma^\infty$, then the subspaces H_σ^2 and \mathcal{N}_σ are invariant for H_Γ^∞ . If $h \in \mathcal{N}_\sigma$, then $fh \in \mathcal{N}_\sigma$ and so $(I - P_{\mathcal{N}_\sigma})M_f P_{\mathcal{N}_\sigma} = 0$. From this we get $(I - P_{\mathcal{N}_\sigma})M_f P_{H_\sigma^2} = (I - P_{\mathcal{N}_\sigma})M_f P_{\mathcal{K}_\sigma}$. If $h \in H_\sigma^2$, then $fh \in H_\sigma^2$, since f is in H_Γ^∞ . Hence, $(I - P_{\mathcal{N}_\sigma})M_f P_{\mathcal{K}_\sigma} = ((I - P_{H_\sigma^2}) + P_{\mathcal{K}_\sigma})M_f P_{\mathcal{K}_\sigma} = P_{\mathcal{K}_\sigma}M_f P_{\mathcal{K}_\sigma}$. Therefore, $\|(I - P_{\mathcal{N}_\sigma})M_f P_{H_\sigma^2}\| = \|P_{\mathcal{K}_\sigma}M_f P_{\mathcal{K}_\sigma}\|$.

Using the fact that $M_f^* k_z^\sigma = \overline{f(z)} k_z^\sigma$ we get that $\|P_{\mathcal{K}_\sigma}M_f P_{\mathcal{K}_\sigma}\| \leq 1$ if and only if the matrices $A_\sigma \geq 0$ for all $\sigma \in \hat{\Gamma}$, where $w_j = f(z_j)$; see [24, Theorem 6.3].

Assume first that $f \in H_\Gamma^\infty$, with $\|f\|_\infty \leq 1$ and $f(z_j) = w_j$. Then $\|P_{\mathcal{K}_\sigma}M_f P_{\mathcal{K}_\sigma}\| \leq \|f\|_\infty \leq 1$ for all $\sigma \in \hat{\Gamma}$.

For the converse, assume that $A_\sigma \geq 0$ for all $\sigma \in \hat{\Gamma}$. By applying [26, Lemma 5.8] we see that there exists a function $g \in H_\Gamma^\infty$ such that $g(z_j) = w_j$. Using the distance formula in Theorem 6.1 we get that $\|g + \mathcal{I}\| = \sup_{\sigma \in \hat{\Gamma}} \|P_{\mathcal{K}_\sigma}M_f P_{\mathcal{K}_\sigma}\| \leq 1$.

Therefore, for each $n \in \mathbb{N}$, there exists $h_n \in \mathcal{I}$ such that $\|g + h_n\|_\infty \leq 1 + n^{-1} \leq 2$. Since the unit ball of H_Γ^∞ is weak* compact, the bounded sequence $f_n := g + h_n$ has a weak* convergent subnet. Let f be the limit of this subnet. Since a point evaluation at a point in the disk is weak* continuous, it follows that $\|f\|_\infty \leq 1$ and $f(z_j) = w_j$. \square

Abrahamse's original theorem is now the special case where Γ is a group of deck transformations that arise from a covering map of a multiply connected domain.

7. TWO EXAMPLES

7.1. Amenable admissible groups. In this section we consider two specific examples of admissible Fuchsian groups. Our purpose with these examples is to illustrate the concepts outlined in this paper. Our first example arises as the group of deck transformations for a covering map of an annulus \mathbb{A} . The group in our second example is isomorphic to $\mathbb{Z}_2 * \mathbb{Z}_2$ and the associated fixed-point algebra is the set of "even" functions in $H^\infty(\mathbb{A})$. Note that both \mathbb{Z} and $\mathbb{Z}_2 * \mathbb{Z}_2$ are amenable. If Γ is an amenable group and s is an invariant mean on $\ell^2(\Gamma)$ we can construct a contractive, unital projection $\Phi_s : H^\infty \rightarrow H_\Gamma^\infty$. We will use this fact later on. On the other hand if Γ is not amenable, then a result of Barrett [10] shows that no such projection can exist.

The first example can also be found in Abrahamse [3, Page 298] and Sarason [28]. Let \mathbb{A} be the annulus with inner radius r and outer radius R . The strip $\mathcal{S} := \{z : 0 < \Re(z) < \pi\}$ is easily seen to be a covering space for the annulus. The covering map is given by $E(z) = \exp\left(\frac{z}{\pi} \log\left(\frac{r}{R}\right)\right)$. The disk and strip are conformally equivalent and we see that the corresponding group of deck transformations is an infinite cyclic group, which we denote Γ_1 , generated by $\gamma(z) = \frac{z-a}{1-az}$, where $0 < a < 1$.

The second example is generated by $\beta(z) = -z$ and $\gamma(z) = \frac{a-z}{1-az}$ with $a \in (0, 1)$. Denote by Γ_2 the group generated by β and γ . Every element of Γ_2 can be identified with a word in β and γ . Since β and γ have order 2, the elements of Γ_2 are words in β and γ with the property that every word is an alternating string of β s and γ s. Hence, there are precisely two words of each length, distinguished by the "letter" they begin with. Set $\alpha = \beta\gamma$. Note that $\alpha(z) = \frac{z-a}{1-az}$ and so Γ_2 contains Γ_1 . A word in Γ_2 is of the form α^m or $\alpha^m\beta$ where $m \in \mathbb{Z}$. Since $\beta(0) = 0$ we see that $\alpha^m(0) = \alpha^m\beta(0)$. If $\alpha^m(0) = 0$, then Lemma 7.1 shows $m = 0$ and so α^m is not the identity map for $m \neq 0$. It follows that $\alpha^m\beta$ is also not the identity map for $m \neq 0$. We have shown that $\Gamma_2 \cong \mathbb{Z}_2 * \mathbb{Z}_2$, the free product of \mathbb{Z}_2 with itself, which is an amenable group.

For these two examples, it can be shown directly that the spaces H_Γ^∞ are non-trivial, that the associated Blaschke product converges, and that we can compute the corresponding character σ_0 . Note that the elements of $H_{\Gamma_2}^\infty$ are exactly the functions in $H_{\Gamma_1}^\infty$ that satisfy $f(z) = f(\beta(z)) = f(-z)$. We will call such a function *even*. The proof of the following lemma involves an elementary induction argument.

Lemma 7.1. Let $\gamma(z) = \frac{z-a}{1-az}$ where $a \in (-1, 1)$ and set

$$a_n = \frac{(1+a)^n - (1-a)^n}{(1-a)^n + (1+a)^n}$$

for $n \geq 1$. We have $\gamma^{(n)}(z) = \frac{z-a_n}{1-a_n z}$ and $\gamma^{(-n)}(z) = \frac{z+a_n}{1+a_n z}$ for $n \geq 1$.

From Lemma 7.1 we see that Γ_1 is admissible.

Lemma 7.2. The Blaschke sum $\sum_{\gamma \in \Gamma_1} (1 - |\gamma(0)|)$ converges.

Proof. We have that the elements of Γ_1 are of the form $\gamma^{(n)}$ where $\gamma(z) = \frac{z-a}{1-az}$ with $a \in (0, 1)$ and $n \in \mathbb{Z}$. From the previous lemma we have

$$1 - a_n = \frac{2(1-a)^n}{(1+a)^n + (1-a)^n} \leq 2 \left(\frac{1-a}{1+a} \right)^n.$$

The latter series is geometric and so the above sum converges. \square

One consequence of this result is that the Blaschke product with zero set $\gamma^{(n)}(0)$ is convergent. Our next proposition gives us σ_0 .

Proposition 7.3. Let B denote the Blaschke product associated to the pair $(\Gamma_1, 0)$. Then $B \circ \gamma = -B$ and $B \circ \beta = -B$. In particular, $B^2 \in H_{\Gamma_2}^\infty$ and the algebra $H_{\Gamma_2}^\infty$ is nontrivial.

Proof. The Blaschke product for the orbit of 0 under Γ_1 is the Blaschke product for the set $\{0, a_n, -a_n : n \geq 1\}$. For $n \in \mathbb{Z}$, let ϕ_n denote the simple Blaschke factor at $\gamma^{(n)}(0)$. For $n \geq 1$, the simple Blaschke factor at $a_n = \gamma^{(-n)}(0)$ is given by

$$\frac{|a_n|}{a_n} \frac{a_n - z}{1 - a_n z} = -\frac{z - a_n}{1 - a_n z} = -\gamma^{(n)}(z).$$

The factor at $-a_n = \gamma^{(n)}(0)$ can be computed similarly and is

$$\frac{|-a_n|}{-a_n} \frac{-a_n - z}{1 + a_n z} = \frac{z + a_n}{1 + a_n z} = \gamma^{(-n)}(z).$$

The factor at 0 is z . Our calculation shows that

$$\phi_n = \begin{cases} \gamma^{(-n)} & n \geq 1 \\ z & n = 0 \\ -\gamma^{(-n)} & n \leq -1 \end{cases}.$$

Let

$$B_N := \prod_{|j| \leq N} \phi_j = (-1)^N \prod_{|j| \leq N} \gamma^{(j)}.$$

Composing with γ and multiplying by $\gamma^{(-N)}$ we get

$$\begin{aligned} \gamma^{(-N)}(B_N \circ \gamma) &= \gamma^{(-N)}(-1)^N \prod_{|j| \leq N} \gamma^{(j+1)} \\ &= ((-1)^N \prod_{|j| \leq N} \gamma^{(j)}) \gamma^{(N+1)} = B_N \gamma^{(N+1)}. \end{aligned}$$

Hence, $\gamma^{(-N)}(z)B_N(\gamma(z)) = \gamma^{(n+1)}(z)B_N(z)$. Since $a_n \rightarrow 1$ as $n \rightarrow \infty$ we get that $\gamma^{(n)}(z) \rightarrow -1$ and $\gamma^{(-n)}(z) \rightarrow 1$ as $n \rightarrow \infty$. Hence, on taking the limit in N we get that $B(\gamma(z)) = -B(z)$.

For $n \in \mathbb{Z}$ we have $\gamma^{(n)}(-z) = -\gamma^{(-n)}(z)$. Hence, $B_N(-z) = (-1)^{2N+1}B_N(z) = -B_N(z)$ and on taking the limit we see $B(-z) = -B(z)$. \square

The stabilizer at the origin for the group Γ_2 consists of the identity and the automorphism β . Hence, by Lemma 2.7, the Blaschke product associated with the pair $(\Gamma_2, 0)$ is B^2 . It follows from Proposition 2.8 that $\{B^{2n} : n \geq 0\}$ is an orthonormal basis for $H_{\Gamma_2}^2$.

7.2. Interpolation results. We now turn our attention to a different approach to the interpolation problem that is valid for the special cases considered in this section. We will deal with the matrix-valued interpolation problem. In the matrix-valued problem we are given n points $z_1, \dots, z_n \in \mathbb{D}$ and n matrices $W_1, \dots, W_n \in M_m$. We seek a function $F \in M_m(H_\Gamma^\infty)$ such that $\|F\|_\infty \leq 1$ and $F(z_j) = W_j$, for $j = 1, \dots, n$.

Since B^2 is inner, we see that $H_{\Gamma_2}^\infty$ is really the span of the powers of an inner function. The interpolation theory for spaces generated by a single inner function is quite simple as Proposition 7.6 will show. To prove Proposition 7.6 we require a few preliminary results.

Lemma 7.4. *If ϕ is an inner function, then $\overline{\phi(\mathbb{D})} = \overline{\mathbb{D}}$.*

Proof. The operator of multiplication by ϕ on H^2 is isometric but not unitary. By the Wold decomposition the spectrum of M_ϕ is the closed unit disk. Hence, $\overline{\mathbb{D}} = \sigma(M_\phi) = \overline{\phi(\mathbb{D})}$. \square

Corollary 7.5. *If ϕ is an inner function and $f \in H^\infty$, then $\|f\|_\infty = \|f \circ \phi\|_\infty$.*

Proof. Since $\phi(\mathbb{D}) \subseteq \mathbb{D}$, $\|f \circ \phi\|_\infty \leq \|f\|_\infty$. If $z \in \mathbb{D}$, then there exists $\phi(z_n) \in \phi(\mathbb{D})$ such that $\phi(z_n) \rightarrow z$ and so $|f(z)| = \lim_{n \rightarrow \infty} |f(\phi(z_n))| \leq \|f \circ \phi\|_\infty$. \square

Proposition 7.6. *Let ϕ be an inner function such that $\phi(0) = 0$. Let H_ϕ^2 be the closed span in H^2 of $\{1, \phi, \phi^2, \dots\}$ and let H_ϕ^∞ be the weak* closure of $\{\phi^n\}_{n \geq 0}$ in H^∞ . We have the following:*

- (1) *The set $\{\phi^n : n \geq 0\}$ is an orthonormal basis for H_ϕ^2 .*
- (2) *The kernel function for H_ϕ^2 is $K^\phi(z, w) = \frac{1}{1 - \phi(z)\overline{\phi(w)}}$.*
- (3) *The function $f \in H_\phi^\infty$ if and only if $f \in H_\phi^2 \cap H^\infty$ if and only if $f = g \circ \phi$ for some $g \in H^\infty$ with $\|g\|_\infty = \|f\|_\infty$.*
- (4) *The multiplier algebra of H_ϕ^2 is H_ϕ^∞ .*
- (5) *The space H_ϕ^2 is a complete Nevanlinna-Pick space.*

Proof. (1) Since the function ϕ vanishes at the origin we have that ϕ^n is orthonormal and since any element of H_ϕ^2 can be approximated by a finite linear combination of the form $a_0 + a_1\phi + \dots + a_n\phi^n$ we see that it is a basis.

(2) We have

$$K^\phi(z, w) = \sum_{n=0}^{\infty} \phi(z)^n \overline{\phi(w)^n} = \frac{1}{1 - \phi(z)\overline{\phi(w)}}$$

- (3) Let us denote by $H^\infty \circ \phi$ the space of functions of the form $\{g \circ \phi : g \in H^\infty\}$ and by \mathcal{A} the weak* closed subalgebra of H^∞ spanned by $\{\phi^n : n \geq 0\}$. The algebra $H^\infty \circ \phi$ is weak* closed and contains \mathcal{A} . If $f \in H_\phi^2 \cap H^\infty$, then $f = \sum_{n=0}^\infty a_n \phi^n$ and so $f = (\sum_{n=0}^\infty a_n z^n) \circ \phi$. The sequence a_n is square summable and so $g = \sum_{n=0}^\infty a_n z^n \in H^2$ and $g \circ \phi = f$. However, by Corollary 7.5 we get that $\|g\|_\infty = \|f\|_\infty$ and so $g \in H^\infty$. This shows that $H_\phi^2 \cap H^\infty \subseteq H^\infty \circ \phi$. If $f \in H^\infty \circ \phi$, then $f = g \circ \phi$ and we can choose a net of polynomials p_t such that $p_t \rightarrow g$ in the weak* topology. Composition by ϕ is weak* continuous and so $p_t \circ \phi \rightarrow f$ in the weak* topology and $\mathcal{A} = H^\infty \circ \phi$. Finally note, since H_ϕ^2 is closed in H^2 , that the space $H_\phi^2 \cap H^\infty$ is weak* closed in H^∞ and contains ϕ^n for all n . This proves that $\mathcal{A} \subseteq H_\phi^2 \cap H^\infty$.
- (4) Since $1 \in H_\phi^2$ and multipliers must be bounded we see that $\text{mult}(H_\phi^2) \subseteq H_\phi^2 \cap H^\infty = H_\phi^\infty$. On the other hand, it is clear from $H_\phi^\infty = H^\infty \circ \phi$ that $H_\phi^\infty \subseteq \text{mult}(H_\phi^2)$. The equality of norms follows as in Proposition 5.1.
- (5) To see that K^ϕ is a complete Nevanlinna-Pick kernel let us consider n points z_1, \dots, z_n in the disk and n matrices W_1, \dots, W_n in M_k . Suppose that the matrix $[(I - W_i W_j^*) K^\phi(z_i, z_j)]_{i,j=1}^n \geq 0$. There exists $F \in M_k(H^\infty)$ such that $\|F\|_\infty \leq 1$ and $F(\phi(z_j)) = W_j$ by the classical Nevanlinna-Pick theorem. The function $\tilde{F} = F \circ \phi$ has the same norm as F and is in H_ϕ^∞ . Also $\tilde{F}(z_j) = F(\phi(z_j)) = W_j$.

On the other hand, suppose that there is a function F in $M_k(H_\phi^\infty)$ such that $F(z_j) = W_j$. We know that $F = G \circ \phi$ for some $G \in M_k(H^\infty)$ with $\|G\|_\infty = \|F\|_\infty$. Hence, $G(\phi(z_j)) = W_j$ and the corresponding Pick matrix is positive. □

Although the proof of Proposition 7.6 is elementary we point out that the representation obtained in part (2) implies the result in part (5) by the work of Agler-McCarthy, McCullough, and Quiggin on complete Nevanlinna-Pick kernels; see Agler and McCarthy's book [5] for a full description.

Applying Proposition 7.6 to the space $H_{\Gamma_2}^2$ we get the following matrix-valued interpolation result.

Corollary 7.7. *Given n points $z_1, \dots, z_n \in \mathbb{D}$ and n matrices $W_1, \dots, W_n \in M_k$, there exists a function $F \in M_k(H_{\Gamma_2}^\infty)$ with $\|F\|_\infty \leq 1$ and $F(z_j) = W_j$ if and only if*

$$\left[\frac{I - W_i W_j^*}{1 - B(z_i)^2 \overline{B(z_j)^2}} \right]_{i,j=1}^n = [(I - W_i W_j^*) K^{\Gamma_2}(z_i, z_j)]_{i,j=1}^n \geq 0,$$

where B is the Blaschke product for the orbit $\Gamma_2(0) = \Gamma_1(0)$.

It is interesting to note that the space $H_{\Gamma_1}^\infty \cong H^\infty(\mathbb{A})$ contains a subalgebra, $H_{\Gamma_2}^\infty$, of "index 2", that is also the multiplier algebra of a complete Nevanlinna-Pick space.

The groups Γ_1 and Γ_2 are amenable. We now take advantage of this fact to study the interpolation problem. Essentially we will project solutions to the classical Nevanlinna-Pick problem onto solutions in H_Γ^∞ .

We briefly discuss group actions on Banach spaces. An action of a group Γ on a Banach space X is a homomorphism from Γ into the group of linear isomorphisms of X . We denote by X_Γ the set of fixed points, that is, $X_\Gamma := \{x \in X : \gamma(x) = x \text{ for all } \gamma \in \Gamma\}$. An action on X induces a natural action on the dual space X^* by $\gamma(f)(x) = f(\gamma(x))$, where $x \in X, f \in X^*$. Conversely, any action of Γ on X^* , where the maps $\gamma : X^* \rightarrow X^*$ are weak* continuous, is induced by an action of Γ on X .

A group Γ is called *amenable* if and only if there exist a linear functional $s : \ell^\infty(\Gamma) \rightarrow \mathbb{C}$ such that $s(1) = 1$, $\|s\| \leq 1$ and s is translation invariant, that is, $s((a_\gamma)_{\gamma \in \Gamma}) = s((a_{\beta^{-1}\gamma})_{\gamma \in \Gamma}) = s((a_{\gamma\beta})_{\gamma \in \Gamma})$ for $\beta \in \Gamma$. Such a linear functional is called an *invariant mean*. In the case where Γ is an amenable group acting on a dual Banach space, there is a natural way to project elements of X^* onto the subspace of fixed-points $(X_\Gamma)^*$. To see this let $x \in X, f \in X^*$ and note that the sequence $\tilde{f} = (f(\gamma(x)))_{\gamma \in \Gamma} \in \ell^\infty(\Gamma)$. Now define $\Phi(f) \in X^*$ by $\Phi(f)(x) = s(\tilde{f})$. From the definition we can check that Φ is a linear, contractive, idempotent map whose range is $(X^*)_\Gamma$ and $\Phi(\gamma^*(f)) = \Phi(f)$ for all $\gamma \in \Gamma$.

The case we are interested in is $X = L^\infty$ and the groups Γ_1 and Γ_2 . In this case Φ is a unital positive map and is therefore completely contractive, that is, given $F = [f_{i,j}]_{i,j=1}^m \in M_m(L^\infty)$, we have $\|\Phi(f_{i,j})\|_{i,j=1}^m \leq \|f_{i,j}\|_{i,j=1}^m$. It follows from the fact that the action of a Fuchsian group restricts to H^∞ that the above projection Φ actually maps H^∞ to H^∞_Γ .

In some ways this new projection is an improvement on the one in [17], since it avoids the defect space N , preserves analytic structure and is completely contractive. The following result shows, in the case of H^∞ , that Φ is in some sense the natural projection to consider.

Proposition 7.8. *Let $f \in H^\infty$ and let the power series expansion of $f \circ \gamma$ be given by $\sum_{n=0}^\infty a_n(\gamma)z^n$. For each n , $(a_n(\gamma))_{\gamma \in \Gamma} \in \ell^\infty(\Gamma)$. If $b_n = s((a_n(\gamma))_{\gamma \in \Gamma})$, then*

$$\Phi(f)(z) = s((f(\gamma(z)))_{\gamma \in \Gamma}) = \sum_{n=0}^\infty b_n z^n.$$

Proof. If k_z is the Szegő kernel at z , then

$$\Gamma(f)(z) = \langle \Phi(f), k_z \rangle = s(\langle f \circ \gamma, k_z \rangle) = s((f(\gamma(z)))_{\gamma \in \Gamma}).$$

We have, $|a_n(\gamma)| = |\langle f \circ \gamma, \chi_n \rangle| \leq \|f\|_\infty$ and so for fixed n , $(a_n(\gamma))_{\gamma \in \Gamma} \in \ell^\infty(\Gamma)$. Now,

$$\langle \Phi(f), z^n \rangle = s(\langle f \circ \gamma, z^n \rangle) = s((a_n(\gamma))_{\gamma \in \Gamma}) = b_n.$$

and so $\Phi(f)(z) = \sum_{n=0}^\infty b_n z^n$. \square

For the special of H^∞ , there is a second averaging result that yields the same projection.

Theorem 7.9. *Let $\mathcal{A} \subseteq B(\mathcal{H})$ and suppose that $\mathcal{A} = \mathcal{A}'$. If Γ is an amenable group that acts on \mathcal{A} , then there exists a linear, contractive, idempotent map $\Phi : \mathcal{A} \rightarrow \mathcal{A}_\Gamma$. If \mathcal{A} is selfadjoint, then so is Φ .*

Proof. Let $A \in \mathcal{A}$ and $f_1, f_2 \in \mathcal{H}$. Define a map $s_A : H \times H \rightarrow \mathbb{C}$ by $s_A(f_1, f_2) := s(\langle \gamma(A)f_1, f_2 \rangle)_{\gamma \in \Gamma}$ and note this is bounded sesquilinear form on \mathcal{H} and that

$\|s_A\| \leq \|A\|$. Therefore, there exists a unique operator $\Phi(A) \in B(H)$ such that $\|\Phi(A)\| = \|s_A\|$ and $\langle \Phi(A)f_1, f_2 \rangle = s(\langle \gamma(A)f_1, f_2 \rangle_{\gamma \in \Gamma})$. Let $A \in \mathcal{A}$ and note that

$$\begin{aligned} \langle \Phi(A)Tf_1, f_2 \rangle &= s(\langle \gamma(A)Tf_1, f_2 \rangle_{\gamma \in \Gamma}) = s(\langle T\gamma(A)f_1, f_2 \rangle_{\gamma \in \Gamma}) \\ &= s(\langle \gamma(A)f_1, T^*f_2 \rangle_{\gamma \in \Gamma}) = \langle \Phi(A)f_1, T^*f_2 \rangle \\ &= \langle T\Phi(A)f_1, f_2 \rangle. \end{aligned}$$

Thus, $\Phi(A) \in \mathcal{A}' = \mathcal{A}$. The remaining properties are easy to verify. \square

It is easy to check, in the case $\mathcal{H} = H^2$ and $\mathcal{A} = H^\infty$, that this agrees with the projection Φ defined earlier.

Proposition 7.10. *If Φ_j is the projection from H^∞ onto $H_{\Gamma_j}^\infty$ for $j = 1, 2$, then $\Phi_j(A(\mathbb{D}))$ is the set of constant functions.*

Proof. Let $\gamma = \frac{z-a}{1-az}$ be as in Lemma 7.1. Since $\gamma^{(k)}(z) \rightarrow -1$ as $k \rightarrow \infty$ and 1 as $k \rightarrow -\infty$ we have that,

$$s(\langle (\gamma^{(k)}(z))^n \rangle_{k \in \mathbb{Z}}) = \begin{cases} 0 & \text{if } k \text{ is odd} \\ 1 & \text{if } k \text{ is even} \end{cases}$$

Therefore, Φ_j maps z^n to a constant and so $A(\mathbb{D})$ is also mapped to the constant functions. \square

Corollary 7.11. *There are no nonconstant, Γ_j -invariant functions in $A(\mathbb{D})$ for $j = 1, 2$.*

Proof. If $f \in A(\mathbb{D})$ and is fixed by Φ_j , then $f = \Phi_j(f)$ is constant. \square

Corollary 7.12. *The projection Φ_j is not weak* continuous.*

Proof. This is a consequence of the fact that the trigonometric polynomials are weak* dense in H^∞ and that $H_{\Gamma_j}^\infty$ contains nonconstant functions. \square

Since interpolation is an isometric theory the existence of a contractive projection suggests, at least in the case of amenable groups, that we could approach the problem of interpolation in H_Γ^∞ through a related problem on the disk. We will show that while this approach has the positive aspect of providing a fairly simple matrix positivity condition, the condition does not seem refined enough to distinguish the interpolation theory of $H_{\Gamma_1}^\infty$ and $H_{\Gamma_2}^\infty$.

Let K denote the Szegő kernel on the disk. By a *weak solution* to the interpolation problem we mean a matrix-valued function $F = [f_{i,j}] \in M_m(H^\infty)$ such that $\|F\|_\infty \leq 1$, $F(\alpha(z_l)) = W_l$ for all $l = 1, \dots, n$ and $\gamma \in \Gamma$. By a *strong solution* we mean an $F = [f_{i,j}] \in M_m(H_\Gamma^\infty)$ such that $\|F\|_\infty \leq 1$ and $F(z_l) = W_l$ for $l = 1, \dots, n$. Clearly every strong solution is a weak solution. If we denote by \mathcal{W} the set of weak solutions, then it is clear, provided that $\mathcal{W} \neq \emptyset$, that \mathcal{W} is a convex, weak*-compact subset of the unit ball of $M_m(H^\infty)$.

Theorem 7.13. *Let Γ be an amenable, admissible group. If $F \in M_m(H^\infty)$ is weak solution, then $\Phi_m(F)$ is a strong solution.*

Proof. We have $\|\Phi_m(F)\| \leq \|F\| \leq 1$, $\Phi_m(F) \in M_m(H_\Gamma^\infty)$ and

$$\Phi_m(F)(z_l) = (\mu(f_{i,j}(\alpha(z_l)))) = W_l.$$

\square

If $Q = [q_{i,j}]_{i,j \in J}$ is an infinite matrix, then we write $Q \geq 0$ to mean that every finite square submatrix is positive. This is also known as *formal positivity*, but we will suppress the word formal. If Q is already finite, or is the matrix of positive operator on a Hilbert space, then the two notions of positivity coincide. The content of the next result is well known and the proof is a standard weak* limit argument.

Proposition 7.14. *Let $\{z_j\}_{j \in J}$ be a set of points in \mathbb{D} and $\{W_j\}_{j \in J} \subseteq M_m$. There exists $F \in M_m(H^{\infty})$ such that $\|F\|_{\infty} \leq 1$ and $f(z_j) = W_j$ if and only the matrix $[(I_m - W_i W_j^*)K(z_i, z_j)]_{i,j \in J} \geq 0$*

If we take the set of points in Proposition 7.14 to be the orbits of n points z_1, \dots, z_n under the group Γ_j , then we obtain the following result:

Corollary 7.15. *Let Γ be an amenable, admissible group. Let $P(z, w)$ denote the infinite matrix $[K(\gamma(z), \eta(w))]_{\gamma, \eta \in \Gamma_j}$. A weak (and hence strong) solution $F \in H_{\Gamma}^{\infty}$ exists if and only if $[(I_m - W_i W_j^*)P(z_i, z_j)]_{i,j=1}^N \geq 0$*

We know that the condition in Corollary 7.15 is equivalent to the condition set forth by Abrahamse for the annulus \mathbb{A} . It would be interesting to have a direct proof that the matrix condition in Corollary 7.15 is equivalent to Abrahamse's condition that the matrices A_{λ} , $\lambda \in \mathbb{T}$ are all positive semi-definite.

Our examples in this section were primarily designed for illustrative purposes. However, we believe that more constructions of this sort could shed further light on the interpolation problem.

We close with an observation that relates to the work of McCullough and Paulsen [23]. Their results show that the C^* -envelope of the n -dimensional quotient algebra $H^{\infty}(\mathbb{A})/\mathcal{I}$ is $M_n(C(\mathbb{T}))$, whenever $n \geq 3$. This should be contrasted with the interpolation result in Corollary 7.7 which shows that the C^* -envelope for the algebra $H_{\Gamma_2}^{\infty}/\mathcal{I}$ is M_n . It was also shown in [11] that the C^* -envelope for any two-idempotent operator algebra [25] is either \mathbb{C}^2 or M_2 . In light of this, three aspects of the interpolation problem, in apparently increasing order of difficulty, would seem to be worthy of study. First, the matrix-valued interpolation problem. This would seem to be within reach using the methods outlined in this paper. Second, a more careful examination of the number of characters required to guarantee the existence of a solution to the interpolation problem. Third, a computation of the C^* -envelope of the quotient algebra $H_{\Gamma}^{\infty}/\mathcal{I}$. In view of the depth of the work in [15] and [23], and in view of the results presented here that these latter problems are considerably more challenging.

REFERENCES

- [1] M. B. Abrahamse, *The Pick interpolation theorem for finitely connected domains.*, Michigan Math. J. **26** (1979), no. 2, 195–203.
- [2] M. B. Abrahamse, *Toeplitz operators in multiply connected regions.*, Amer. J. Math., **96** (1974), 261–297. MR0361891
- [3] M. B. Abrahamse, *Analytic Toeplitz operators with automorphic symbol.*, Proc. Amer. Math. Soc., **52** (1975), 297–302. MR0405156
- [4] M. B. Abrahamse and R. G. Douglas, *A class of subnormal operators related to multiply-connected domains.*, Advances in Math., **19** (1976), no. 1, 106–148.
- [5] Jim Agler and John E. McCarthy, *Pick interpolation and Hilbert function spaces.*, Graduate Studies in Mathematics, 44. American Mathematical Society, Providence, RI, 2002. MR1882259

- [6] P. R. Ahern and Donald Sarason, *The H^p spaces of a class of function algebras.*, Acta Math., **117** (1967), 123–163.
- [7] W.B. Arveson, *Interpolation problems in nest algebras*, J. Funct. Anal., **20** (1975), 208–233. MR0383098
- [8] Joseph A. Ball, *Operator extremal problems, expectation operators and applications to operators on multiply connected domains.*, J. Operator Theory, **1** (1979), no. 2, 153–175. MR0532872
- [9] Joseph A. Ball and Victor Vinnikov, *Hardy spaces on a finite bordered Riemann surface, multivariable operator model theory and Fourier analysis along a unimodular curve*. Systems, approximation, singular integral operators, and related topics (Bordeaux, 2000), 37–56, Oper. Theory Adv. Appl., 129, Birkhuser, Basel, 2001. MR1882690
- [10] David E. Barrett, *Failure of averaging on multiply connected domains.*, Ann. Inst. Fourier (Grenoble), **40** (1990), no. 2, 357–370. MR1070831
- [11] Kenneth R. Davidson, Vern I. Paulsen, Mrinal Raghupathi and Dinesh Singh, *A constrained Nevanlinna-Pick interpolation problem*, Indiana Univ. Math. J., to appear.
- [12] C. J. Earle and A. Marden, *On Poincaré series with application to H^p spaces on bordered Riemann surfaces.*, Illinois J. Math., **13** (1969), 202–219. MR0237766
- [13] Ronald G. Douglas, *Banach algebra techniques in operator theory.*, Second edition. Graduate Texts in Mathematics, 179. Springer-Verlag, New York, 1998. MR1634900
- [14] Hershel M. Farkas and Irwin Kra, *Riemann surfaces.*, Second edition. Graduate Texts in Mathematics, 71. Springer-Verlag, New York, 1992. MR1139765
- [15] V. L. Vinnikov and S. I. Fedorov, *On the Nevanlinna-Pick interpolation in multiply connected domains.*, translation in J. Math. Sci. (New York), **105** (2001), no. 4, 2109–2126. MR1691394
- [16] Stephen D. Fisher, *Function theory on planar domains. A second course in complex analysis*. John Wiley & Sons, Inc., New York, 1983. MR0694693
- [17] Frank Forelli, *Bounded holomorphic functions and projections.*, Illinois J. Math., **10** (1966), 367–380. MR0193534
- [18] Morisuke Hasumi, *Invariant subspace theorems for finite Riemann surfaces.*, Canad. J. Math., **18** (1966), 240–255. MR0190790
- [19] Henry Helson, *Harmonic analysis.*, Second edition. Hindustan Book Agency, Delhi, 1995. MR1373194
- [20] Kenneth Hoffman, *Banach spaces of analytic functions*. Dover Publications, Inc., New York, 1988. MR1102893
- [21] Svetlana Katok, *Fuchsian groups*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1992. MR1177168
- [22] Scott McCullough, *Isometric representations of some quotients of H^∞ of an annulus*. Integral Equations Operator Theory, **39** (2001), no. 3, 335–362. MR1818066
- [23] Scott McCullough and Vern Paulsen, *C^* -envelopes and interpolation theory.*, Indiana Univ. Math. J., **51** (2002), no. 2, 479–505. MR1909298
- [24] Vern I. Paulsen, An introduction to the theory of reproducing kernel Hilbert space, course notes, available at <http://www.math.uh.edu/~vern>
- [25] Vern I. Paulsen, *Operator algebras of idempotents.*, J. Funct. Anal., **181** (2001), no. 2, 209–226. MR1821697
- [26] Mrinal Raghupathi, *Nevanlinna-Pick interpolation for $\mathbb{C}+BH^\infty$* , Integral Equation Operator Theory (2009), to appear.
- [27] Donald Sarason, *Generalized interpolation in H^∞ .*, Trans. Amer. Math. Soc., **127** (1967), 179–203. MR0208383
- [28] Donald Sarason, *The H^p spaces of an annulus.*, Mem. Amer. Math. Soc., **56** (1965). MR0188824
- [29] Joel H. Shapiro, *Composition operators and classical function theory*. Universitext: Tracts in Mathematics. Springer-Verlag, New York, 1993. MR1237406
- [30] M. Tsuji, *Potential theory in modern function theory*. Chelsea Publishing Co., New York, 1975. MR0414898
- [31] Michael Voichick, *Ideals and invariant subspaces of analytic functions.*, Trans. Amer. Math. Soc., **111** (1964), 493–512. MR0160920
- [32] Michael Voichick and Lawrence Zalcman, *Inner and outer functions on Riemann surfaces.*, Proc. Amer. Math. Soc., **16** (1965), 1200–1204. MR0183883

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TENNESSEE, 37240,
U.S.A.

E-mail address: `mrinal.raghupathi@vanderbilt.edu`

URL: `http://www.math.vanderbilt.edu/people/raghupathi`